

**2026 1 Tier Standard - SCHA**  
**2026 Prior Authorization Criteria**  
 CURRENT AS OF 05/01/2026

# ACITRETIN

---

**Products Affected**

- *acitretin*

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescriber must be a dermatologist or an oncologist.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	For prophylaxis of skin cancer in patients with previously treated skin cancers who have undergone an organ transplantation the request will be approved. For psoriasis: the patient has documented trial of, contraindication to, or medical reason for not using at least 2 of the treatment options listed: topical steroids, tazarotene, methotrexate, and cyclosporine.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# ACL INHIBITORS

## Products Affected

- NEXLETOL
- NEXLIZET

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	18 years or older
<b>Prescriber Restrictions</b>	Prescriber must be a cardiologist or specialist in the treatment of lipid disorders
<b>Coverage Duration</b>	New starts will be authorized for 4 months. Cont of therapy or reauth until end of contract year.
<b>Other Criteria</b>	For new starts ALL of the following must be provided: 1) Documentation of baseline low density lipoprotein cholesterol (LDL-C), 2) Member has tried and failed a high-intensity statin (i.e. atorvastatin 40-80 mg, rosuvastatin 20-40 mg) at maximum tolerated dose for 3 months via claim history or chart notes OR documentation has been provided that the member is not able to tolerate a statin. In addition to the initial criteria above if the new start is for the diagnosis of hypercholesterolemia, the following are required: 1) Member has a diagnosis of heterozygous familial hypercholesterolemia (FH) OR hypercholesterolemia, 2) Member has tried ezetimibe at a maximum tolerated dose and LDL-C is not at goal, or documentation has been provided that the patient is not able to tolerate ezetimibe. In addition to the initial criteria above if the new start is for cardiovascular risk reduction, the following are required: 1) Member has established cardiovascular disease (documented history of coronary artery disease, symptomatic peripheral artery disease, and/or cerebrovascular atherosclerotic disease, 2) Member does not have established cardiovascular disease but is considered high risk (one of the following): Diabetes Mellitus (Type 1 or Type 2) in females over 65 years of age or males over 60 years of age OR a Reynolds Risk score greater than 30% or a SCORE Risk score greater than 7.5% over 10 years OR a coronary artery calcium score greater than 400 Agaston units at any time in the past, 3) Member has a fasting LDL-C greater than or equal to 70 mg/dL. For

<b>PA Criteria</b>	<b>Criteria Details</b>
	continuation of therapy or reauthorization requests for all indications: Documentation provided that the member has obtained clinical benefit from medication (e.g., LDL-C lowering from baseline)
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# ACTEMRA

## Products Affected

- ACTEMRA ACTPEN
- ACTEMRA SUBCUTANEOUS

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Specialist for submitted diagnosis.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	For pJIA: Either 1) Trial of, medical reason for not using, or contraindication to 2 of the following: Enbrel, an adalimumab product, or Xeljanz or 2) If utilized within the past 120 days, approve for continuation of therapy. For RA: Either 1) Trial of, medical reason for not using, or contraindication to 2 of the following: Enbrel, an adalimumab product, or Xeljanz or 2) If utilized within the past 120 days, approve for continuation of therapy. For sJIA, Giant Cell Arteritis and Systemic Sclerosis-Associated Interstitial Lung Disease: Approve
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# ADALIMUMAB

## Products Affected

- *adalimumab-fkjp (2 pen)*
- *adalimumab-fkjp (2 syringe) subcutaneous prefilled syringe kit 20 mg/0.4ml, 40 mg/0.8ml*
- SIMLANDI (1 PEN) SUBCUTANEOUS AUTO-INJECTOR KIT 40 MG/0.4ML, 80 MG/0.8ML
- SIMLANDI (1 SYRINGE)
- SIMLANDI (2 PEN)
- SIMLANDI (2 SYRINGE) SUBCUTANEOUS PREFILLED SYRINGE KIT 20 MG/0.2ML, 40 MG/0.4ML

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Specialist for submitted diagnosis.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	For ankylosing spondylitis: Trial of, medical reason for not using, or contraindication to naproxen. For pJIA: Trial of, medical reason for not using, or contraindication to 1 of the following DMARDs: methotrexate or leflunomide. For RA: Trial of, medical reason for not using, or contraindication to 1 disease modifying antirheumatic drug (DMARD) (methotrexate, leflunomide, or sulfasalazine). For PsA, psoriasis, Hidradenitis Suppurativa, Crohn's Disease (CD), Ulcerative Colitis (UC) or Uveitis: approve.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Prerequisite Therapy Required</b>	Yes

# ADEMPAS

## Products Affected

- ADEMPAS

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Concurrent use with Phosphodiesterase Inhibitors used for Pulmonary Hypertension or Other Soluble Guanylate Cyclase Stimulators.
<b>Required Medical Information</b>	Pulmonary arterial hypertension (PAH) (World Health Organization [WHO] Group 1)- Initial: [Note: documentation required] (1) PAH was confirmed by right heart catheterization AND (2) pretreatment mean pulmonary arterial pressure is greater than 20 mmHg AND (3) pretreatment pulmonary capillary wedge pressure is less than or equal to 15 mmHg, AND (4) pretreatment pulmonary vascular resistance is greater than or equal to 3 Wood units. Continuation of therapy: patient has positive clinical response to treatment. Chronic thromboembolic pulmonary hypertension (CTEPH) (WHO Group 4) - Initial: [Note: documentation required] (1) Patient has persistent or recurrent CTEPH after pulmonary endarterectomy (PEA) OR (2) patient has inoperable CTEPH with the diagnosis confirmed by right heart catheterization AND by computed tomography (CT), magnetic resonance imaging (MRI), or pulmonary angiography. Continuation of therapy: patient has positive clinical response to treatment.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a cardiologist or pulmonologist.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

Formulary ID 26312  
Last Update: 04/21/2026

# ALPHA-1 PROTEINASE INHIBITORS

---

## Products Affected

- ARALAST NP INTRAVENOUS SOLUTION RECONSTITUTED 1000 MG, 500 MG
- GLASSIA
- PROLASTIN-C INTRAVENOUS SOLUTION
- ZEMAIRA

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Documentation of hereditary alpha1-antitrypsin deficiency as evident by pretreatment serum AAT levels below 11 micromol/L and progressive FEV1 or FVC decline demonstrating symptomatic lung disease.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescriber must be a pulmonologist.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	If the medication request is for Glassia or Aralast NP, the patient has a documented medical reason (such as trial, intolerance or contraindication) for not using Prolastin-C or Zemaira to treat their medical condition.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# ALYFTREK

---

## Products Affected

- ALYFTREK ORAL TABLET 10-50-125 MG, 4-20-50 MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	Concurrent use with Trikafta, Kalydeco, Orkambi, or Symdeko. Patients with unknown CFTR gene mutations.
<b>Required Medical Information</b>	Documentation of CFTR gene that is responsive to vanzacaftor-tezacaftor-deutivacaftor treatment.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a pulmonologist or a provider who specializes in treatment of CF.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	Cystic Fibrosis (CF) - Initial: patient must have at least one F508del mutation in the CFTR gene or a mutation in the CFTR gene that is responsive to the requested medication. Continuation of therapy: approve.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# AMBRISENTAN

---

## Products Affected

- *ambrisentan*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Pulmonary arterial hypertension (PAH) (World Health Organization [WHO] Group 1)- Initial: [Note: documentation required] (1) PAH was confirmed by right heart catheterization AND (2) pretreatment mean pulmonary arterial pressure is greater than 20 mmHg AND (3) pretreatment pulmonary capillary wedge pressure is less than or equal to 15 mmHg, AND (4) pretreatment pulmonary vascular resistance is greater than or equal to 3 Wood units. Continuation of therapy: patient has positive clinical response to treatment.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a cardiologist or pulmonologist.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# ANTINEOPLASTIC AGENTS

---

## Products Affected

- *abiraterone acetate*
- ABIRTEGA
- AKEEGA
- ALECENSA
- ALUNBRIG
- AUGTYRO
- AVMAPKI FAKZYNJA CO-PACK
- AYVAKIT
- BALVERSA
- *bexarotene*
- BOSULIF
- BRAFTOVI ORAL CAPSULE 75 MG
- BRUKINSA
- CABOMETYX
- CALQUENCE ORAL TABLET
- CAPRELSA
- COMETRIQ (100 MG DAILY DOSE) ORAL KIT 80 & 20 MG
- COMETRIQ (140 MG DAILY DOSE) ORAL KIT 3 X 20 MG & 80 MG
- COMETRIQ (60 MG DAILY DOSE)
- COPIKTRA
- COTELLIC
- DANZITEN
- *dasatinib*
- DAURISMO
- ENSACOVE
- ERIVEDGE
- ERLEADA
- *erlotinib hcl*
- EULEXIN
- *everolimus oral tablet 10 mg, 2.5 mg, 5 mg, 7.5 mg*
- *everolimus oral tablet soluble*
- FOTIVDA
- FRUZAQLA
- GAVRETO
- *gefitinib*
- GILOTRIF
- GOMEKLI
- HERNEXEOS
- HYRNUO
- IBRANCE
- IBTROZI
- ICLUSIG
- IDHIFA
- *imatinib mesylate oral*
- IMKELDI
- INLURIYO
- INLYTA
- INQOVI
- INREBIC
- ITOVEBI
- IWILFIN
- JAYPIRCA
- KISQALI (200 MG DOSE)
- KISQALI (400 MG DOSE)
- KISQALI (600 MG DOSE)
- KISQALI FEMARA (400 MG DOSE)
- KISQALI FEMARA (600 MG DOSE)
- KOMZIFTI
- KOSELUGO
- KRAZATI
- *lapatinib ditosylate*
- LAZCLUZE
- *lenalidomide*
- LENVIMA (10 MG DAILY DOSE)
- LENVIMA (12 MG DAILY DOSE)
- LENVIMA (14 MG DAILY DOSE)
- LENVIMA (18 MG DAILY DOSE)
- LENVIMA (20 MG DAILY DOSE)
- LENVIMA (24 MG DAILY DOSE)
- LENVIMA (4 MG DAILY DOSE)
- LENVIMA (8 MG DAILY DOSE)
- LEUKERAN
- LONSURF
- LORBRENA
- LUMAKRAS
- LYNPARZA ORAL TABLET
- LYTGobi (12 MG DAILY DOSE)
- LYTGobi (16 MG DAILY DOSE)
- LYTGobi (20 MG DAILY DOSE)
- MEKINIST
- MEKTOVI

Formulary ID 26312

Last Update: 04/21/2026

- *mercaptopurine oral suspension*
- MODEYSO
- NERLYNX
- *nilotinib d-tartrate*
- *nilotinib hcl*
- *nilutamide*
- NINLARO
- NUBEQA
- ODOMZO
- OGSIVEO ORAL TABLET 100 MG, 150 MG
- OJEMDA
- OJJAARA
- ONUREG
- ORGOVYX
- ORSERDU
- *pazopanib hcl*
- PEMAZYRE
- PIQRAY (200 MG DAILY DOSE)
- PIQRAY (250 MG DAILY DOSE)
- PIQRAY (300 MG DAILY DOSE)
- *pomalidomide*
- QINLOCK
- RETEVMO ORAL TABLET
- REVLIMID
- REVUFORJ
- REZLIDHIA
- ROMVIMZA
- ROZLYTREK
- RUBRACA
- RYDAPT
- SCEMBLIX
- SOLTAMOX
- *sorafenib tosylate*
- STIVARGA
- *sunitinib malate*
- TABLOID
- TABRECTA
- TAFINLAR
- TAGRISSO
- TALZENNA
- TAZVERIK
- TEPMETKO
- THALOMID ORAL CAPSULE 100 MG, 50 MG
- TIBSOVO
- *toremifene citrate*
- *tretinoin oral*
- TRUQAP ORAL TABLET 200 MG
- TRUQAP ORAL TABLET THERAPY PACK
- TUKYSA
- TURALIO ORAL CAPSULE 125 MG
- VANFLYTA
- VENCLEXTA
- VENCLEXTA STARTING PACK
- VERZENIO
- VITRAKVI
- VIZIMPRO
- VONJO
- VORANIGO
- WELIREG
- XALKORI
- XOSPATA
- XPOVIO (100 MG ONCE WEEKLY) ORAL TABLET THERAPY PACK 50 MG
- XPOVIO (40 MG ONCE WEEKLY) ORAL TABLET THERAPY PACK 10 MG, 40 MG
- XPOVIO (40 MG TWICE WEEKLY) ORAL TABLET THERAPY PACK 40 MG
- XPOVIO (60 MG ONCE WEEKLY) ORAL TABLET THERAPY PACK 60 MG
- XPOVIO (60 MG TWICE WEEKLY)
- XPOVIO (80 MG ONCE WEEKLY) ORAL TABLET THERAPY PACK 40 MG, 80 MG
- XPOVIO (80 MG TWICE WEEKLY)
- XTANDI
- YONSA
- ZEJULA ORAL TABLET
- ZELBORAF
- ZOLINZA
- ZYDELIG
- ZYKADIA ORAL TABLET

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescriber must be an oncologist or specialist for submitted diagnosis.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# APOMORPHINE

---

## Products Affected

- *apomorphine hcl subcutaneous*

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	Concomitant use with serotonin 5-HT3 receptor antagonists.
<b>Required Medical Information</b>	Reviewer will verify available patient claim history to confirm patient is not using 5-HT3 receptor antagonists.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a neurologist.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	If diagnosis is Parkinson's, the patient must have a documented trial of, contraindication to, or medical reason for not using two alternatives such as entacapone, tolcapone, rasagiline, selegiline, carbidopa/levodopa, bromocriptine, pramipexole or ropinirole.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# AQNEURSA

---

## Products Affected

- AQNEURSA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Specialist for submitted diagnosis.
<b>Coverage Duration</b>	New starts will be authorized for 6 months. Cont of therapy or reauth until end of contract year.
<b>Other Criteria</b>	For new starts: 1) The member has a documented diagnosis of Niemann-Pick disease type C (NPC) AND 2) Documentation of genetic testing identifying disease-causing alleles in NPC1 or NPC2 AND 3) Documentation of disease-related neurological symptoms (e.g., developmental delay/regression, ataxia, cataplexy, seizures, motor-function decline, tremors, dysphagia) For reauthorization: Documentation that member has had positive response to therapy (e.g., stabilization in neurological status, decrease in functional Scale for Assessment and Rating of Ataxia [fSARA] score).
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# ARCALYST

---

## Products Affected

- ARCALYST

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Specialist for submitted diagnosis.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	For deficiency of interleukin-1 receptor antagonist, documented trial of, contraindication to, or medical reason for not using Kineret. For continuation of therapy or reauthorization: Documentation has been provided that patient has clinically benefited from medication.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# ARIKAYCE

---

## Products Affected

- ARIKAYCE

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Mycobacterium avium complex (MAC): (1) Documented diagnosis of MAC lung disease as verified by failure to achieve at least 2 negative sputum cultures following 6 consecutive months of a combination antibacterial drug regimen AND (2) Provider attestation that medication is being used as part of a combination antibacterial drug regimen.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a pulmonologist or an infectious disease specialist
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# ARISTADA

## Products Affected

- ARISTADA INITIO 441 MG/1.6ML, 662 MG/2.4ML, 882
- ARISTADA INTRAMUSCULAR MG/3.2ML
- PREFILLED SYRINGE 1064 MG/3.9ML,

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	The member has a documented history of receiving oral aripiprazole without any clinically significant side effects.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	Trial and failure of, contraindication, or medical reason (e.g. intolerance, hypersensitivity or contraindication) for not using at least two of the following: Abilify Maintena, Abilify Asimtufii, Invega Sustenna, Invega Trinza, Invega Hafyera, or Risperidone Microspheres ER.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# AUVELITY

---

## Products Affected

- AUVELITY

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	Seizure disorder
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	Trial of, contraindication to, or medical reason for not using two generic antidepressants.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# AZTREONAM LYSINE

---

## Products Affected

- CAYSTON

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescriber must be a pulmonologist, infectious disease specialist, or an expert in the treatment of cystic fibrosis.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# BENLYSTA

---

## Products Affected

- BENLYSTA SUBCUTANEOUS

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescriber must be a rheumatologist, nephrologist, or specialist in the treatment of autoimmune disorders.
<b>Coverage Duration</b>	New starts will be authorized for 6 months. Cont of therapy or reauth until end of contract year.
<b>Other Criteria</b>	For new starts for systemic lupus erythematosus (SLE): concurrent use of two of the following or medical reason for not using glucocorticoids, azathioprine, methotrexate, mycophenolate, or hydroxychloroquine, chloroquine, and cyclophosphamide. For continuation of therapy or reauthorization for SLE: documentation of clinical response to therapy (i.e. fewer flares that required steroid treatment, lower average daily oral prednisone dose, improved daily function either as measured through a validated functional scale or through improved daily performance documented at clinic visits, etc.) For new starts for lupus nephritis (LN): concurrent use of or medical reason for not using background immunosuppressive therapy regimen. For continuation of therapy or reauthorization for LN: Documentation of improvement in renal function (i.e. reduction in UPCR).
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Prerequisite Therapy Required</b>	No

# BESREMI

---

## Products Affected

- BESREMI

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescriber must be a hematologist, oncologist, or specialist for submitted diagnosis.
<b>Coverage Duration</b>	The request will be authorized until the end of the contract year.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# BOSENTAN

---

## Products Affected

- *bosentan oral tablet*

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Documentation of pulmonary arterial hypertension (PAH) WHO Group I classification and PAH Functional Class.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescriber must be a pulmonologist or cardiologist.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# BRINSUPRI

---

## Products Affected

- BRINSUPRI

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	Patients with bronchiectasis due to cystic fibrosis.
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	12 years of age and older.
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a pulmonologist.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	For non-cystic fibrosis bronchiectasis - Initial: (1) patient has documented history of bronchiectasis diagnosed by chest computed tomography scan AND (2) prescriber confirmation that bronchiectasis is not primarily driven by chronic obstructive pulmonary disease or asthma. Continuation of therapy: patient has positive clinical response to treatment.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# CAMZYOS

## Products Affected

- CAMZYOS

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a cardiologist.
<b>Coverage Duration</b>	New starts will be authorized for 6 months. Cont of therapy or reauth until end of contract year.
<b>Other Criteria</b>	For all new starts, documentation of ALL of the following must be provided: 1) Diagnosis of symptomatic New York Heart Association (NYHA) class II or III obstructive hypertrophic cardiomyopathy (oHCM) AND 2) Patient has a left ventricular ejection fraction (LVEF) greater than or equal to 55% AND 3) Assessment of Valsalva left ventricular outflow tract (LVOT) gradient AND 4) Trial of, medical reason for not using or contraindication to BOTH of the following: Beta blockers (i.e. metoprolol, propranolol, atenolol) AND Non-dihydropyridine calcium channel blockers (i.e. verapamil, diltiazem) AND 5) Prescriber attests that patient is not using moderate to strong CYP2C19 or CYP3A4 inhibitors or inducers. For continuation of therapy or reauthorization, all of the following must be provided: 1) Documentation of clinical benefit as evidenced by an improvement from baseline in oHCM symptoms (i.e., improvement in fatigue, chest pain, shortness of breath, LVOT, peak oxygen consumption, etc.) OR improvement or no worsening of NYHA functional class AND 2) Member must also have a left ventricular ejection fraction (LVEF) greater than or equal to 50%.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# CARDAMYST

---

## Products Affected

- CARDAMYST

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a cardiologist or specialist for submitted diagnosis.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	Documented history of acute symptomatic episodes of paroxysmal supraventricular tachycardia.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# CARGLUMIC ACID

---

## Products Affected

- *carglumic acid oral tablet soluble*

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# CASPOFUNGIN

---

## Products Affected

- *casposfungin acetate*

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Documentation of a consultation with an infectious disease specialist.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# CERDELGA

---

## Products Affected

- CERDELGA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	Patients with undetermined CYP2D6 metabolizer status.
<b>Required Medical Information</b>	Patient's CYP2D6 metabolizer status, as determined by an FDA approved test. For reauthorization, documentation has been provided that patient has obtained clinical benefit from medication (e.g. increased platelet count, improvement in anemia, PFTs, improvement in radiographic scans, improved quality of life).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a specialist in treatment of Gaucher's disease.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# CGRP ANTAGONISTS

## Products Affected

- AIMOVIG
- EMGALITY
- EMGALITY (300 MG DOSE)
- NURTEC
- QULIPTA
- UBRELVY
- ZAVZPRET

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	New starts will be authorized for 6 months. Cont of therapy or reauth until end of contract year.
<b>Other Criteria</b>	For acute migraine new starts - for Ubrelvy and Nurtec requests, must have trial of, contraindication to or medical reason for not using a triptan. For Zavzpret requests, must have trial of, contraindication to or medical reason for not using Ubrelvy or Nurtec. For migraine prophylaxis new starts - at least 4 migraine days per month or one or more severe migraine attacks lasting for greater than 12 hours despite use of abortive therapy (e.g. triptans or NSAIDs). For Emgality requests for episodic cluster headache new starts - approve. For continuation of therapy or reauthorization - For acute migraine (Nurtec, Ubrelvy, Zavzpret), must show documentation of improvement in migraine symptoms (pain, photophobia, phonophobia). For migraine prevention (Nurtec, Emgality, Qulipta, Aimovig), must show documentation of improvement in migraine symptoms. For episodic cluster headache treatment, must show documentation of reduction in frequency of headaches.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

Formulary ID 26312  
Last Update: 04/21/2026

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Prerequisite Therapy Required</b>	Yes

# CHOLBAM

---

## Products Affected

- CHOLBAM

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	For new starts: Patient has documented diagnosis of either: 1) bile acid synthesis disorder due to a single enzyme defect or 2) peroxisomal disorders. For continuation of therapy or reauthorization: prescriber attests: 1) the patient has clinical improvement with therapy (i.e. liver function tests) AND 2) there is no evidence of biliary obstruction or cholestasis
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescriber must be a hepatologist, gastroenterologist, or metabolic specialist
<b>Coverage Duration</b>	New starts will be authorized for 3 months. Cont of therapy or reauth until end of contract year.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# CIBINQO

---

## Products Affected

- CIBINQO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Specialist for submitted diagnosis.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	For atopic dermatitis: Trial of, contraindication to, or medical reason for not using Rinvoq or Dupixent.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# CIMZIA

## Products Affected

- CIMZIA (1 SYRINGE)
- CIMZIA (2 SYRINGE)
- CIMZIA SUBCUTANEOUS KIT 2 X 200 MG
- CIMZIA-STARTER

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Specialist for submitted diagnosis.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	For ankylosing spondylitis: Either 1) Trial of, medical reason for not using, or contraindication to 2 of the following therapies: Enbrel, an adalimumab product, Cosentyx, or Xeljanz or 2) If utilized within the past 120 days, approve for continuation of therapy. For Crohns Disease: Either 1) Trial of, medical reason for not using, or contraindication to 2 of the following therapies: an adalimumab product or an ustekinumab product or 2) If utilized within the past 120 days, approve for continuation of therapy. For non-radiographic axial spondyloarthritis: approve. For psoriasis: Either 1) Trial of, medical reason for not using, or contraindication to 2 of the following therapies: Tremfya, an ustekinumab product, Enbrel, or an adalimumab product or 2) If utilized within the past 120 days, approve for continuation of therapy. For PsA: Either 1) Trial of, medical reason for not using, or contraindication to 2 of the following therapies: Cosentyx, an ustekinumab product, Tremfya, Xeljanz, Enbrel, or an adalimumab product or 2) If utilized within the past 120 days, approve for continuation of therapy. For RA: Either 1) Trial of, medical reason for not using, or contraindication to 2 of the following: Enbrel, an adalimumab product, or Xeljanz or 2) If utilized within the past 120 days, approve for continuation of therapy. Polyarticular juvenile idiopathic arthritis (pJIA) - Initial: trial of, medical reason for not using, or contraindication to 2 of the following:

<b>PA Criteria</b>	<b>Criteria Details</b>
	Enbrel, an adalimumab product or Xeljanz or or 2) If utilized within the past 120 days, approve for continuation of therapy.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# CORLANOR

---

## Products Affected

- CORLANOR ORAL SOLUTION
- *ivabradine hcl*

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	New starts for chronic heart failure must have all of the following: 1) LVEF of 35% or less 2) Sinus rhythm and have resting heart rate greater than or equal to 70 bpm. For pediatric patients with heart failure due to dilated cardiomyopathy: approve.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescriber must be a cardiologist.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	Trial of, contraindication to, or medical reason for not receiving a beta blocker.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# CORTROPHIN

## Products Affected

- CORTROPHIN
- CORTROPHIN GEL

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	New starts for MS exacerbation, rheumatic disorders, collagen diseases, dermatologic diseases, serum sickness, edematous state (e.g. nephrotic syndrome without uremia), and respiratory diseases: trial of, contraindication to, or medical reason for not using oral corticosteroids. New starts for ophthalmic disease: trial of, contraindication to, or medical reason for not using oral or ophthalmic corticosteroids. Continuation of therapy or reauthorization for MS exacerbation: documentation of symptom improvement and current use of a multiple sclerosis disease modifying agent for maintenance therapy. Continuation of therapy or reauthorization for all other conditions: documented evidence of response to treatment and symptom improvement.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Specialist for submitted diagnosis.
<b>Coverage Duration</b>	MS exacerbation: 1 month. Other conditions: new start for 3 months and reauth end of contract year.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# COSENTYX

---

## Products Affected

- COSENTYX (300 MG DOSE)
- COSENTYX INTRAVENOUS
- COSENTYX SENSOREADY (300 MG)
- COSENTYX SENSOREADY PEN
- COSENTYX SUBCUTANEOUS SOLUTION PREFILLED SYRINGE 150 MG/ML, 75 MG/0.5ML
- COSENTYX UNOREADY

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	1) Documented diagnosis of ankylosing spondylitis, non-radiographic axial spondyloarthritis, plaque psoriasis, psoriatic arthritis, enthesitis-related arthritis, rheumatoid arthritis or hidradenitis suppurativa. 2) If utilized within the past 120 days, approve for continuation of therapy.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No
Prerequisite Therapy Required	No

# CRESEMBA

---

## Products Affected

- CRESEMBA ORAL

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Documented diagnosis of medically accepted indication.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an infectious disease specialist or oncologist
<b>Coverage Duration</b>	Request will be authorized for 3 months.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# CYSTAGON

---

## Products Affected

- CYSTAGON

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# CYSTARAN

---

## Products Affected

- CYSTARAN

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Documentation of diagnosis for cystinosis with corneal cystine crystal accumulation.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an ophthalmologist or metabolic disease specialist.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# DALFAMPRIDINE ER

---

## Products Affected

- *dalfampridine er*

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	History of seizure or moderate/severe renal impairment (CrCl less than or equal to 50 mL/min).
<b>Required Medical Information</b>	For new starts: 1) Attestation that creatinine clearance (CrCl) greater than 50 mL/min was confirmed prior to initiation of therapy, AND 2) Documentation has been provided that member is ambulatory (able to walk at least 25 feet) and has a documented walking impairment, AND 3) For multiple sclerosis, member is currently being treated with a disease modifying agent (e.g. immunomodulator, interferon, etc.) or has a medical reason why member is unable to use a disease modifying agent for their condition. For continuation of therapy or re-authorization requests: 1) Member must experience improvement in walking from baseline due to use of dalfampridine ER.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescriber must be a neurologist.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# DEFERASIROX

## Products Affected

- *deferasirox*
- *deferasirox granules*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Creatinine clearance less than 40 mL/min or platelet counts less than 50,000/mm <sup>3</sup> .
<b>Required Medical Information</b>	For all indications: platelet count greater than or equal to 50,000/mm <sup>3</sup> (within 30 days). For chronic iron overload due to transfusions: serum ferritin concentration greater than 1000 mcg/L (lab result within 30 days). For chronic iron overload in non-transfusion-dependent thalassemia syndromes: serum ferritin concentration greater than 300 mcg/L (lab result within 30 days).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	For deferasirox granules oral packets, the member must have medical reason for not using deferasirox tablets or oral soluble tablets.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# DEFERIPRONE

---

## Products Affected

- *deferiprone*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	For new starts: 1) serum ferritin level above 1,000 mcg/L and absolute neutrophil count (ANC) greater than $1.5 \times 10^9/L$ within 30 days of request, and 2) Trial of, contraindication to, or medical reason for not using deferasirox tablets. For continuation of therapy or reauthorization, decrease in serum ferritin from baseline.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

# DEFLAZACORT

---

## Products Affected

- KYMBEE

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	1) Member has a documented diagnosis of Duchenne Muscular Dystrophy as evidenced by one of the following: a) documented mutation of dystrophin gene OR b) muscle biopsy indicating absence of the dystrophin protein AND 2) Patient has trial and failure with, contraindication to or medical reason for not taking prednisone.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a neurologist or specialist for submitted diagnosis.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# DIACOMIT

---

## Products Affected

- DIACOMIT

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescriber must be a neurologist.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	For members 2 years and older: Trial of, contraindication to, or medical reason for not using one generic anticonvulsant for appropriate indications. For members under 2 years old: Approve.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# DIHYDROERGOTAMINE NASAL

---

## Products Affected

- *dihydroergotamine mesylate nasal*

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	For new starts: Member has a diagnosis of migraine headaches with or without aura. Prescriber attestation that it will be used for the acute treatment of migraine. For continuation of therapy or reauthorization: Documentation or provider attestation of positive clinical response (e.g., improvement in pain, photophobia, phonophobia).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Requests will be authorized for 12 weeks.
<b>Other Criteria</b>	Trial of, contraindication to, or medical reason (e.g. intolerance or hypersensitivity) for not using a triptan (e.g., rizatriptan, sumatriptan).
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# DOPTELET

---

## Products Affected

- DOPTELET
- DOPTELET SPRINKLE

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	For new starts for chronic liver disease and chronic immune thrombocytopenia (chronic ITP): documented baseline platelet count of less than 50,000/mcL.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with hematologist, hepatologist or surgeon.
<b>Coverage Duration</b>	For thrombocytopenia with CLD getting procedure: 5 days. For all others: remainder of contract year
<b>Other Criteria</b>	For chronic ITP: trial of, contraindication to, or medical reason for not using a corticosteroid. For thrombocytopenia with chronic liver disease (CLD): approve.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# DOXEPIN CREAM

---

## Products Affected

- *doxepin hcl external*

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Request will be authorized for 1 month.
<b>Other Criteria</b>	Trial of, contraindication to, or medical reason for not using a topical corticosteroid or topical calcineurin inhibitor.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# DUPIXENT

## Products Affected

- DUPIXENT SUBCUTANEOUS SOLUTION AUTO-INJECTOR
- DUPIXENT SUBCUTANEOUS SOLUTION PREFILLED SYRINGE 200 MG/1.14ML, 300 MG/2ML

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Chronic spontaneous urticaria (CSU) - Initial: patient has inadequate symptomatic relief despite trial of two weeks of a second-generation oral antihistamine (unless contraindicated). Continuation of therapy: patient has positive clinical response to treatment. Bullous pemphigoid - Initial: patient has had trial of, contraindication to, or medical reason for not using a topical corticosteroid, oral corticosteroid, or an immunosuppressive agent. Continuation of therapy: patient has positive clinical response to treatment.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Initial therapy only: prescribed by or in consultation with specialist for submitted diagnosis.
<b>Coverage Duration</b>	PN: Initial 6 mo. Cont of therapy and all others initial: end of contract year.
<b>Other Criteria</b>	Atopic dermatitis (AD) in patients 6 months of age and older - Initial: (1) patient has diagnosis of moderate to severe AD, AND (2) has had trial of, contraindication to, or medical reason for not using either a topical corticosteroid or topical calcineurin inhibitor. Continuation of therapy: patient has positive clinical response to treatment. Asthma with eosinophilic phenotype - Initial: (1) patient has baseline blood eosinophil count greater than or equal to 150 cells per microliter, AND (2) asthma remains inadequately controlled despite current treatment with or medical reasons for not using BOTH (i) medium to high dose inhaled corticosteroid AND (ii) additional controller (i.e. leukotriene modifier, long acting beta-2-agonist, long acting muscarinic antagonist, and sustained released theophylline). Asthma, oral corticosteroid dependent - Initial: asthma remains inadequately controlled despite current treatment with or medical reasons for not using BOTH (i) high dose inhaled corticosteroid AND (ii) additional controller (i.e. leukotriene modifier, long acting beta-2-agonist, long acting muscarinic antagonist, and sustained released theophylline).

PA Criteria	Criteria Details
	<p>Asthma with eosinophilic phenotype or oral corticosteroid dependent - Continuation of therapy: clinical improvement in asthma control (i.e. reduction in frequency and/or severity of exacerbations and symptoms OR reduction in daily maintenance oral corticosteroid dose). Chronic rhinosinusitis with nasal polyps (CRSwNP) - Initial: (1) Dupixent is used as add-on maintenance treatment, AND (2) patient has a trial of, contraindication to, or medical reason for not using nasal corticosteroids. Continuation of therapy: patient has positive clinical response to treatment. Prurigo nodularis (PN): attestation is provided confirming diagnosis. Eosinophilic esophagitis (EoE) - Initial: (1) diagnosis has been confirmed by esophageal biopsy AND (2) patient weighs at least 15 kilograms AND (3) patient experienced an inadequate treatment response, intolerance, or has a contraindication to a topical or oral corticosteroid (i.e. fluticasone propionate or budesonide). Continuation of therapy: patient has positive clinical response to treatment. Chronic Obstructive Pulmonary Disease (COPD) - Initial: (1) documented diagnosis of COPD with an eosinophilic phenotype, AND (2) Dupixent is used as an add-on maintenance treatment, AND (3) documentation that patient's COPD is inadequately controlled. Continuation of therapy: patient has positive clinical response to treatment.</p>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# EGRIFTA

---

## Products Affected

- EGRIFTA SV
- EGRIFTA WR

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Documentation of active antiretroviral therapy for at least 8 weeks.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# EMSAM

---

## Products Affected

- EMSAM

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	Concomitant use with SSRIs, SNRIs, clomipramine and imipramine, meperidine, tramadol, methadone, pentazocine, and propoxyphene, and the antitussive agent dextromethorphan or carbamazepine
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	For new starts: Trial of, contraindication to, or medical reason for not using two generic antidepressants.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# ENBREL

## Products Affected

- ENBREL MINI
- ENBREL SUBCUTANEOUS SOLUTION 25 MG/0.5ML
- ENBREL SUBCUTANEOUS SOLUTION PREFILLED SYRINGE
- ENBREL SURECLICK SUBCUTANEOUS SOLUTION AUTO-INJECTOR

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Specialist for submitted diagnosis.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	For RA: Trial of, medical reason for not using, or contraindication to 1 disease modifying antirheumatic drug (DMARD) (methotrexate, leflunomide, or sulfasalazine). For pJIA: Trial of, medical reason for not using, or contraindication to 1 of the following DMARDs: methotrexate or leflunomide. For PsA or psoriasis: approve. For ankylosing spondylitis: Trial of, medical reason for not using, or contraindication to naproxen. Continuation of therapy: patient has been receiving Enbrel for a minimum of 4 months and has positive clinical response to treatment.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

# ENDARI

---

## Products Affected

- *l-glutamine oral packet*

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Documentation that two or more painful sickle cell crises have occurred in the past 12 months.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescriber must be a hematologist.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	Trial of, contraindication to, or medical reason for not using hydroxyurea for at least three months.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# ENTYVIO

---

## Products Affected

- ENTYVIO PEN

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Specialist for submitted diagnosis.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	For ulcerative colitis: Either 1) Trial of, medical reason for not using, or contraindication to 2 of the following therapies: an ustekinumab product, Tremfya, or an adalimumab product or 2) If utilized within the past 120 days, approve for continuation of therapy.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# EPIDIOLEX

---

## Products Affected

- EPIDIOLEX

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescriber must be a neurologist.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	Trial of, contraindication to, or medical reason for not using one generic anticonvulsant for appropriate indications.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# EPRONTIA

---

## Products Affected

- *topiramate oral solution*

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	The request will be authorized until the end of the contract year.
<b>Other Criteria</b>	Documented trial of, contraindication to, or medical reason for not using topiramate sprinkle capsule or topiramate oral tablet.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# ERYTHROPOIETIN STIMULATING AGENTS

## Products Affected

- ARANESP (ALBUMIN FREE) INJECTION SOLUTION 100 MCG/ML, 200 MCG/ML, 25 MCG/ML, 40 MCG/ML, 60 MCG/ML
- ARANESP (ALBUMIN FREE) INJECTION SOLUTION PREFILLED SYRINGE
- EPOGEN INJECTION SOLUTION 10000 UNIT/ML, 2000 UNIT/ML, 20000 UNIT/ML, 3000 UNIT/ML, 4000 UNIT/ML, 4000 UNIT/ML
- PROCIT
- RETACRIT INJECTION SOLUTION 10000 UNIT/ML, 10000 UNIT/ML(1ML), 2000 UNIT/ML, 20000 UNIT/ML, 3000 UNIT/ML, 4000 UNIT/ML, 40000 UNIT/ML

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	For new starts for all indications: Hgb within compendia range for treatment of the requested medical condition. For continuation of therapy or re-authorization: Hgb must not exceed 10 g/dL (anemia related to cancer), 11 g/dL (anemia of CKD), 12 g/dL (zidovudine-related anemia in members with HIV and ribavirin-induced anemia), 13 g/dL (elective, noncardiac, nonvascular surgery needing red blood cell allogeneic transfusion reduction).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Request will be authorized for 6 months.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

Formulary ID 26312  
Last Update: 04/21/2026



# EUCRISA

---

## Products Affected

- EUCRISA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescriber must be a dermatologist, immunologist or an allergist.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	For patients 2 years of age and older: Trial of, contraindication to, or medical reason for not using topical tacrolimus or pimecrolimus. For patients less than 2 years of age: approve.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# EVRYSDI

---

## Products Affected

- EVRYSDI

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	For new starts, all of the following must be included: 1) Documentation of genetic testing confirming diagnosis AND 2) Documentation of baseline motor function or motor milestone achievement [e.g. CHOP Infant Test of Neuromuscular Disorders (CHOP-INTEND) or Hammersmith Infant Neurological Examination (HINE) for Type 1 or Hammersmith Functional Motor Scale Expanded Scores (HFMSE), or Total Motor Function Measure 32 (MFM-32) for Type II and Type III, or 6 minute walk test in subjects able to walk]. For continuation of therapy or reauthorization, documentation of clinical response has been submitted (e.g. improvement in motor function/motor milestone achievement scores using CHOP-INTEND or HFMSE, MFM-32, 6 minute walk test or HINE improvement in more categories of motor milestones than worsening).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescriber must be a neurologist.
<b>Coverage Duration</b>	New starts will be authorized for 6 months. Cont of therapy or reauth until end of contract year.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# FABHALTA

## Products Affected

- FABHALTA

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Concurrent use with another complement inhibitor for the treatment of PNH (i.e. Empaveli, Soliris, or Ultomiris).
<b>Required Medical Information</b>	PNH - Initial: patient has documented diagnosis of paroxysmal nocturnal hemoglobinuria (PNH) confirmed by (1) flow cytometry analysis confirming presence of PNH clones AND (2) patient has signs and symptoms of PNH (i.e. anemia, abdominal pain, dyspnea, kidney disease, pulmonary hypertension, hemolysis/hemoglobinuria, etc.). Continuation of therapy: patient has documented positive clinical response to treatment (i.e. decrease in LDH, increased or stabilization of hemoglobin levels, reduction in transfusions, increased reticulocyte count, etc.). Reduction of proteinuria in adults with immunoglobulin A (IgA) nephropathy - Initial: patient has documented diagnosis of IgA nephropathy AND IgA nephropathy at risk of rapid disease progression (i.e. clinical evidence of rapid disease progression generally a urine protein-to-creatinine ratio or UPCR greater or equal to 1.5g/g OR other clinically relevant tests). Continuation of therapy: patient has documented positive clinical response to treatment.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a hematologist, nephrologist or oncologist.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

Formulary ID 26312  
Last Update: 04/21/2026

# FASENRA

## Products Affected

- FASENRA
- FASENRA PEN

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Specialist for submitted diagnosis.
<b>Coverage Duration</b>	New starts will be authorized for 6 months. Cont of therapy or reauth until end of contract year.
<b>Other Criteria</b>	New starts for severe asthma with an eosinophilic phenotype: 1)Baseline blood eosinophil count greater than or equal to 150 cells per microliter AND 2) symptoms persist with at least 1 exacerbation in the last 12 months requiring additional treatment (e.g. oral systemic steroids) while on a high dose inhaled corticosteroid with an additional controller medication (ie. long-acting B2 agonist). Continuation of therapy or re-authorization for severe asthma with an eosinophilic phenotype: clinical benefit from use of the drug. New starts for eosinophilic granulomatosis with polyangiitis (EGPA)- Initial: patient has a documented history or the presence of an eosinophil count of more than 1000 cells per microliter or a blood eosinophil level of greater than 10 percent AND trial of, contraindication to, or medical reason for not using one of the following medications: cyclophosphamide or methotrexate. Continuation of therapy: patient has a beneficial response to treatment with the requested drug (i.e. a reduction in the frequency of relapses, decrease in the daily oral corticosteroid dose, or no active vasculitis).
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

Formulary ID 26312  
 Last Update: 04/21/2026

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Prerequisite Therapy Required</b>	Yes

# FILSPARI

---

## Products Affected

- FILSPARI

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	Coadministration with renin-angiotensin-aldosterone system (RAAS) inhibitors, endothelin receptor antagonists, or aliskiren
<b>Required Medical Information</b>	For new starts: Documentation is provided that the member has diagnosis of primary immunoglobulin A nephropathy (IgAN) at risk of rapid disease progression. Member has proteinuria. For continuation of therapy or reauthorization: Documentation of positive clinical response (ie. decrease in urine protein-to-creatinine ratio (UPCR)).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a nephrologist.
<b>Coverage Duration</b>	New starts will be authorized for 9 months. Cont of therapy or reauth until end of contract year.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# FINTEPLA

---

## Products Affected

- FINTEPLA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescriber must be a neurologist.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	Trial of, contraindication to, or medical reason for not using one generic anticonvulsant for appropriate indications.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# FIRDAPSE

---

## Products Affected

- FIRDAPSE

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	History of seizures.
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescriber must be a neurologist.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# FLUCYTOSINE

---

## Products Affected

- *flucytosine oral*

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	Complete dihydropyrimidine dehydrogenase (DPD) enzyme deficiency
<b>Required Medical Information</b>	Attestation member is taking in combination with amphotericin B.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# GALAFOLD

---

## Products Affected

- GALAFOLD

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed must be a geneticist, cardiologist, nephrologist or specialist experienced in the treatment of Fabry disease.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# GATTEX

---

## Products Affected

- GATTEX

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	For new starts: attestation of 1) Colonoscopy and upper gastrointestinal endoscopy with removal of polyps within six months prior to starting treatment for adults or 2) Fecal occult blood testing within six months prior to starting treatment for pediatric patients and 3) Baseline laboratory assessments (bilirubin, alkalinephosphatase, lipase, and amylase). For continuation of therapy or reauthorization: Documentation is provided that the member has obtained a clinical benefit (e.g. reduction in parenteral fluid volume, reduction in number of days receiving parenteral nutrition).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a gastroenterologist.
<b>Coverage Duration</b>	New starts will be authorized for 6 months. Cont of therapy or reauth until end of contract year.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# GLP-1 AGONISTS

---

## Products Affected

- *liraglutide*
- MOUNJARO SUBCUTANEOUS SOLUTION AUTO-INJECTOR
- OZEMPIC (0.25 OR 0.5 MG/DOSE) SUBCUTANEOUS SOLUTION PEN-INJECTOR 2 MG/3ML
- OZEMPIC (1 MG/DOSE) SUBCUTANEOUS SOLUTION PEN-INJECTOR 4 MG/3ML
- OZEMPIC (2 MG/DOSE)
- RYBELSUS
- TRULICITY SUBCUTANEOUS SOLUTION AUTO-INJECTOR

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	The member has an indication of weight loss/obesity only or type 1 diabetes. The member has concurrent use of any GLP-1 receptor agonist.
<b>Required Medical Information</b>	The member has a documented diagnosis via chart notes of type 2 diabetes.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	Diabetes: patient has diagnosis of type 2 diabetes mellitus. All other indications: patient must have medically accepted indication.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# GNRH AGONISTS

---

## Products Affected

- ELIGARD
- FIRMAGON (240 MG DOSE)
- FIRMAGON SUBCUTANEOUS SOLUTION RECONSTITUTED 80 MG
- *leuprolide acetate (3 month)*
- *leuprolide acetate injection*
- LUPRON DEPOT (1-MONTH)
- LUPRON DEPOT (3-MONTH)
- LUPRON DEPOT (4-MONTH)
- LUPRON DEPOT (6-MONTH)
- LUTRATE DEPOT
- TRELSTAR MIXJECT

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Fibroids and endometriosis: 6 months. All other indications: end of contract year.
Other Criteria	If the medication request is for the treatment of prostate cancer and if the request is for any other GnRH agonist other than Eligard or leuprolide, the patient must have a documented trial of, contraindication to, or medical reason for not using Eligard or leuprolide to treat their prostate cancer.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

# GOCOVRI

---

## Products Affected

- GOCOVRI

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a neurologist.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	New starts: trial of, contraindication to, or medical reason for not using generic amantadine. Continuation of therapy or reauthorization: Member demonstrates clinical benefit (i.e. improvement in levodopa-induced dyskinesia or decreased off episodes).
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# GROWTH HORMONES

## Products Affected

- GENOTROPIN MINIQUICK SUBCUTANEOUS PREFILLED SYRINGE
- GENOTROPIN SUBCUTANEOUS CARTRIDGE
- NGENLA
- OMNITROPE SUBCUTANEOUS SOLUTION CARTRIDGE
- OMNITROPE SUBCUTANEOUS SOLUTION RECONSTITUTED
- SKYTROFA

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	For new starts for growth hormone deficiency: Documentation showing bone age testing, height, weight, and Growth Hormone Stimulation Test results OR Insulin Growth Factor 1 level. For continuation of therapy or reauthorization for growth hormone deficiency: documentation (medical records) showing positive response to treatment.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescriber must be an endocrinologist or nephrologist.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	For new starts for growth hormone deficiency: 1) If the request is not for Genotropin, trial of, contraindication to, or medical reason for not using Genotropin. For requests for all other medically accepted indications other than growth hormone deficiency, the request will be approved for products other than Skytrofa.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# HEREDITARY ANGIOEDEMA AGENTS

---

## Products Affected

- CINRYZE
- HAEGARDA
- ORLADEYO

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescriber must be an allergist, immunologist, rheumatologist or hematologist.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	For continuation of therapy or reauthorization: Documentation has been provided that patient has clinically benefited from medication.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No
Prerequisite Therapy Required	No

# HIGH RISK MEDICATION

## Products Affected

- *benztropine mesylate oral*
- *cyproheptadine hcl oral*
- *diphenoxylate-atropine oral liquid*
- *diphenoxylate-atropine oral tablet 2.5-0.025 mg*
- *dipyridamole oral*
- *ergotamine-caffeine*
- *glyburide micronized oral tablet 3 mg, 6 mg*
- *glyburide oral tablet 1.25 mg, 2.5 mg, 5 mg*
- *glyburide-metformin oral tablet 1.25-250 mg*
- *glyburide-metformin oral tablet 2.5-500 mg, 5-500 mg*
- *guanfacine hcl er oral tablet extended release 24 hour 1 mg, 2 mg, 3 mg, 4 mg*
- *guanfacine hcl oral*
- *hydroxyzine hcl oral syrup 10 mg/5ml*
- *hydroxyzine hcl oral tablet*
- *hydroxyzine pamoate oral*
- *indomethacin oral capsule 25 mg, 50 mg*
- *ketorolac tromethamine oral*
- *nifedipine oral*
- *promethazine hcl oral solution 6.25 mg/5ml*
- *promethazine hcl oral tablet*
- *promethazine hcl rectal suppository 12.5 mg, 25 mg*
- *promethazine-phenylephrine*
- *promethegan rectal suppository 50 mg*

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	For patients 65 years old and older the prescriber has documented: 1) the benefits of treatment with the drug outweigh the potential risks and 2) the risks and side effects have been discussed and will be monitored.
<b>Age Restrictions</b>	Prior authorization only applies to members 65 years old or older.
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

Formulary ID 26312  
Last Update: 04/21/2026

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Prerequisite Therapy Required</b>	No

# HIGH RISK MEDICATION - PROTECTED CLASS DRUGS

## Products Affected

- *amitriptyline hcl oral*
- *amoxapine*
- *clomipramine hcl oral*
- *nortriptyline hcl oral*
- *perphenazine-amitriptyline*
- *phenobarbital oral elixir*
- *phenobarbital oral tablet*

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	For patients 65 years old and older the prescriber has documented: 1) the benefits of treatment with the drug outweigh the potential risks and 2) the risks and side effects have been discussed and will be monitored.
<b>Age Restrictions</b>	Prior authorization only applies to members 65 years old or older.
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# HIGH RISK MEDICATION, BUTALBITAL

## Products Affected

- *bac (butalbital-acetamin-caff)*
- *butalbital-acetaminophen oral tablet 50-325 mg*
- *butalbital-apap-caff-cod oral capsule 50-325-40-30 mg*
- *butalbital-apap-caffeine oral capsule 50-325-40 mg*
- *butalbital-apap-caffeine oral solution*
- *butalbital-apap-caffeine oral tablet 50-325-40 mg*
- *butalbital-asa-caff-codeine*
- *butalbital-aspirin-caffeine oral capsule*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	For patients 65 years old and older the prescriber has documented: 1) the benefits of treatment with the drug outweigh the potential risks and 2) the risks and side effects have been discussed and will be monitored.
<b>Age Restrictions</b>	Prior authorization only applies to members 65 years old or older.
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	Trial of, contraindication to, or medical reason for not using an oral NSAID.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# HIGH RISK MEDICATION, MEGESTROL

## Products Affected

- *megestrol acetate oral suspension 40 mg/ml, 400 mg/10ml, 625 mg/5ml*
- *megestrol acetate oral tablet*

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	For patients 65 years old and older the prescriber has documented: the benefits of treatment with the drug outweigh the potential risks identified for people 65 years old and older.
<b>Age Restrictions</b>	Prior authorization only applies to members 65 years old or older.
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# HIGH RISK MEDICATION, SHORT TERM MUSCLE RELAXANT

## Products Affected

- *carisoprodol oral*
- *chlorzoxazone oral tablet 500 mg*
- *cyclobenzaprine hcl oral tablet 10 mg, 5 mg*
- *metaxalone oral tablet 800 mg*
- *methocarbamol oral tablet 500 mg, 750 mg*
- *orphenadrine citrate er*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	For patients 65 years old and older the prescriber has documented: 1) the benefits of treatment with the drug outweigh the potential risks identified for people 65 years old and older, and 2) the risks and side effects have been discussed and will be monitored.
<b>Age Restrictions</b>	Prior authorization only applies to members 65 years old or older.
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	New starts will be authorized for 30 days. Continuation of therapy or reauth will be for 90 days.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# HIGH RISK MEDICATION, SLEEP AGENTS

## Products Affected

- *eszopiclone*
- *temazepam*
- *zaleplon oral capsule 10 mg, 5 mg*
- *zolpidem tartrate er*
- *zolpidem tartrate oral tablet 10 mg*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	For patients 65 years old and older the prescriber has documented: 1) the benefits of treatment with the drug outweigh the potential risks identified for people 65 years old and older, and 2) the risks and side effects have been discussed and will be monitored. For zolpidem immediate release 10mg and zolpidem ER: trial of or medical reason for not using zolpidem immediate release 5mg.
<b>Age Restrictions</b>	Prior authorization only applies to members 65 years old or older.
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# HYFTOR

---

## Products Affected

- HYFTOR

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	For new starts: documentation of diagnosis of tuberous sclerosis with facial angiofibroma. For continuation of therapy or reauthorization: documentation that the member has experienced a clinical benefit from treatment (e.g. improvement in size and color of angiofibroma).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a neurologist or provider who specializes in the treatment of genetic or dermatologic disorders.
<b>Coverage Duration</b>	New starts: 3 months. Cont. of therapy or reauthorization: until end of contract year.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# ICATIBANT

---

## Products Affected

- *icatibant acetate subcutaneous solution prefilled syringe*

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescriber must be an immunologist, allergist, rheumatologist, or hematologist.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# ILARIS

---

## Products Affected

- ILARIS SUBCUTANEOUS SOLUTION

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Documentation was submitted indicating that the member was evaluated for active or latent TB infection (i.e. tuberculin skin test)
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Specialist for submitted diagnosis.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	For sJIA: approve.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# ILUMYA

---

## Products Affected

- ILUMYA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Specialist for submitted diagnosis.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	For psoriasis: Either 1) Trial of, medical reason for not using, or contraindication to 2 of the following therapies: an ustekinumab product, Tremfya, Enbrel, or an adalimumab product or 2) If utilized within the past 120 days, approve for continuation of therapy.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# IMBRUVICA

---

## Products Affected

- IMBRUVICA ORAL CAPSULE
- IMBRUVICA ORAL SUSPENSION
- IMBRUVICA ORAL TABLET 140 MG, 280 MG, 420 MG

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescriber must be an oncologist or specialist for submitted diagnosis.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	For new starts for treatment of graft-versus-host disease (GVHD): Trial of, contraindication to, or medical reason for not using a systemic corticosteroid. For continuation of therapy of for treatment of GVHD: documentation of clinical benefit from use of the drug (i.e. symptom improvement, reduction in corticosteroid dose). For all other indications, approve.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# IMPAVIDO

---

## Products Affected

- IMPAVIDO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Documentation of diagnosis with one of the following: (a) Visceral leishmaniasis due to <i>Leishmania donovani</i> , (b) Cutaneous leishmaniasis due to <i>Leishmania braziliensis</i> , <i>Leishmania guyanensis</i> , or <i>Leishmania panamensis</i> , (c) Mucosal leishmaniasis due to <i>Leishmania braziliensis</i> .
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Request will be authorized for 28 days.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# INCRELEX

---

## Products Affected

- INCRELEX

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an endocrinologist.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# JAKAFI

---

## Products Affected

- JAKAFI

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescriber must be an oncologist or specialist for submitted diagnosis.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	For new starts for treatment of graft-versus-host disease (GVHD): Trial of, contraindication to, or medical reason for not using a systemic corticosteroid. For continuation of therapy of for treatment of GVHD: documentation of clinical benefit from use of the drug (i.e. symptom improvement, reduction in corticosteroid dose). For all other indications, approve.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# KALYDECO

---

## Products Affected

- KALYDECO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	Combination use with Orkambi, Symdeko, or Trikafta.
<b>Required Medical Information</b>	Documentation of CFTR gene that is responsive to ivacaftor treatment.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescriber must be a pulmonologist or an expert in the treatment of cystic fibrosis.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# KERENDIA

## Products Affected

- KERENDIA

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	For new starts: 1) Documentation of diagnosis of chronic kidney disease (CKD) due to type 2 diabetes mellitus OR heart failure (HF) with left ventricular ejection fraction (LVEF) greater than or equal to 40% AND AND 2) Documentation of serum potassium levels less than or equal to 5 mEq/L AND 3) eGFR greater than or equal to 25ml/min/1.73 m <sup>2</sup> AND 4) Documentation that member is taking Kerendia in combination with an angiotensin-converting enzyme inhibitor (ACEi) or angiotensin receptor blocker (ARB) at maximum tolerated doses or documentation has been provided that the member is unable to tolerate ACEi or ARB. For continuation of therapy or reauthorization for patients with CKD due to type 2 diabetes mellitus: 1) Documentation of serum potassium levels less than or equal to 5.5 mEq/L AND 2) Documentation that member is taking Kerendia in combination with an ACEi or ARB at maximum tolerated doses or documentation has been provided that the member is unable to tolerate ACEi or ARB. For continuation of therapy or reauthorization for patients with HF with LVEF greater than or equal to 40%: 1) Documentation of serum potassium levels less than 6 mEq/L AND 2) Documentation that member is taking Kerendia in combination with an ACEi or ARB at maximum tolerated doses or documentation has been provided that the member is unable to tolerate ACEi or ARB.
<b>Indications</b>	All Medically-accepted Indications.

Formulary ID 26312

Last Update: 04/21/2026

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# KEVZARA

---

## Products Affected

- KEVZARA

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Specialist for submitted diagnosis.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	For RA: Either 1) Trial of, medical reason for not using, or contraindication to 2 of the following: Enbrel, an adalimumab product, or Xeljanz or 2) If utilized within the past 120 days, approve for continuation of therapy. For polymyalgia rheumatica (PMR): Trial of, medical reason for not using, or contraindication to corticosteroids. For pJIA: Either 1) Trial of, medical reason for not using, or contraindication to 2 of the following: Enbrel, an adalimumab product, or Xeljanz or 2) If utilized within the past 120 days, approve for continuation of therapy.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# KINERET

---

## Products Affected

- KINERET SUBCUTANEOUS SOLUTION PREFILLED SYRINGE

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Specialist for submitted diagnosis.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	For RA: Either 1) Trial of, medical reason for not using, or contraindication to 2 of the following: Enbrel, an adalimumab product, or Xeljanz or 2) If utilized within the past 120 days, approve for continuation of therapy. For cryopyrin-associated periodic syndromes or deficiency of interleukin-1 receptor antagonist: Approve.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# LEQEMBI IQLIK

## Products Affected

- LEQEMBI IQLIK

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescriber must be a neurologist, geriatric psychiatrist, or geriatrician who specializes in treating dementia
<b>Coverage Duration</b>	Initial: 6 months. Continuation of therapy: end of the contract year.
<b>Other Criteria</b>	Initiation: (1) the patient has documentation of 18 months of treatment with intravenous Leqembi prior to initiation of subcutaneous maintenance therapy & (2) the patient has a diagnosis of mild cognitive impairment (MCI) caused by Alzheimer's Disease (AD) or mild AD consistent with Stage 3 or Stage 4 Alzheimer's disease & (3) documentation of both of the following: a positive result for the presence of beta-amyloid plaques & baseline Magnetic Resonance Imaging (MRI) scan. Continuation of therapy: patient has a positive clinical response to treatment.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# LEQSELVI

---

## Products Affected

- LEQSELVI

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Documentation was submitted indicating that the member was evaluated for active or latent TB infection (i.e. tuberculin skin test).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	Documentation of confirmed diagnosis and other causes of hair loss have been ruled out.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# LITFULO

---

## Products Affected

- LITFULO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Initial: (1) documented diagnosis via chart notes of severe alopecia areata AND (2) patient is not receiving in combination with either of the following: (i) Targeted immunomodulator (i.e. Olumiant, Enbrel, Cimzia, Simponi, Orencia, adalimumab, Xeljanz, Rinvoq) OR (ii) potent immunosuppressant. Continuation of therapy: documentation of positive clinical response to treatment.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	Documentation of confirmed diagnosis and other causes of hair loss have been ruled out.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# LIVMARLI

## Products Affected

- LIVMARLI

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a gastroenterologist or hepatologist.
<b>Coverage Duration</b>	New starts will be authorized for 6 months. Cont of therapy or reauth until end of contract year.
<b>Other Criteria</b>	For new starts: 1) Trial of, contraindication to, or medical reason for not using both of the following: cholestyramine AND rifampin. 2) Prescriber attests that the member has cholestasis 3) Baseline serum bile acid level is provided. 4) Documentation of patients weight. For continuation of therapy or reauthorization: 1) Documentation submitted indicating the member has had all of the following: an improvement in pruritis (e.g. improved observed scratching, decreased sleep disturbances/nighttime awakenings due to scratching, etc.) AND reduction in serum bile acid level from baseline. 2) Prescriber attests that patient has had no evidence of hepatic decompensation (e.g. variceal hemorrhage, ascites, hepatic encephalopathy, portal hypertension, etc.). 3) Documentation of patients weight.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# LIVTENCITY

---

## Products Affected

- LIVTENCITY

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Cytomegalovirus (CMV): (1) Documented diagnosis of CMV infection AND (2) Member is a recipient of one of the following: (a) hematopoietic stem cell transplant, (b) solid organ transplant AND (3) patient has tried and failed treatment with valganciclovir, ganciclovir, cidofovir, or foscarnet AND (4) patient weighs greater than or equal to 35 kg.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a transplant, oncologist, or infectious disease specialist.
<b>Coverage Duration</b>	Request will be authorized for 8 weeks.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# LODOCO

---

## Products Affected

- LODOCO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescriber must be, or in consultation with a specialist in the treatment of cardiovascular disease, such as a cardiologist
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	Documentation that patient has established atherosclerotic disease or multiple risk factors for cardiovascular disease AND documentation that patient does not have pre-existing blood dyscrasias (ex. leukopenia, thrombocytopenia) and patient does not have renal failure (CrCl less than 15 ml/min) or severe hepatic impairment
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# LUCEMYRA

---

## Products Affected

- *lofexidine hcl*

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Request will be authorized for 14 days.
<b>Other Criteria</b>	Patient must have trial of, contraindication to, or medical reason for not using clonidine.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# LUPKYNIS

## Products Affected

- LUPKYNIS

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Concurrent use with cyclophosphamide.
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescriber must be rheumatologist, nephrologist, or other specialist in the treatment of autoimmune disorders.
<b>Coverage Duration</b>	New starts will be authorized for 6 months. Cont of therapy or reauth until end of contract year.
<b>Other Criteria</b>	For new starts: 1) Documentation of urine protein/creatinine ratio (UPCR), 2) Documentation that the member has a baseline eGFR greater than 45 mL/min/1.73m <sup>2</sup> or that benefit outweighs risk of using this medication at current eGFR, and 3) Concurrent use of or medical reason for not using background immunosuppressive therapy regimen. For continuation of therapy or reauthorization: Documentation of improvement in renal function (i.e. reduction in UPCR or no confirmed decrease from baseline eGFR greater than or equal to 20%).
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# LYBALVI

---

## Products Affected

- LYBALVI

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	Concomitant use with opioids.
<b>Required Medical Information</b>	Attestation from the provider that the member has had an opioid-free period of a minimum of 7 days after last use of shorting-acting opioids and 14 days from last use of long-acting opioids before initiating Lybalvi.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	Documented trial of, contraindication to, or medical reason for not using at least two generic antipsychotics, one of which must be generic olanzapine.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# MAVYRET

---

## Products Affected

- MAVYRET

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Detectable HCV RNA viral load prior to treatment within 6 months of request. In addition, documentation of treatment history, and if cirrhotic, documentation of compensated or decompensated cirrhosis.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Request will be authorized for 8-16 weeks as per AASLD-IDSA guidance.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# METHOTREXATE ORAL SOLUTION

---

## Products Affected

- JYLAMVO
- XATMEP

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescriber must be an oncologist, a rheumatologist, a dermatologist, or other appropriate specialist
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	Polyarticular Juvenile Idiopathic Arthritis (pJIA) - Initial: (1) patient had been diagnosed with pJIA AND (2) patient had tried, intolerant or has medical reason for not using at least one non-steroidal anti-inflammatory agents (NSAIDs) AND methotrexate.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# METHYLTESTOSTERONE

---

## Products Affected

- *methyltestosterone oral*

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# METYROSINE

---

## Products Affected

- *metyrosine*

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Documentation of one of the following: 1) Concurrent use of alpha adrenergic blockers, 2) Medical reason for being unable to use an alpha adrenergic blocker, OR 3) Patient is not a candidate for surgical resection and requires long term treatment with metyrosine.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# MIFEPRISTONE

---

## Products Affected

- *mifepristone oral tablet 300 mg*

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	For all members patient must not be currently on simvastatin, lovastatin, cyclosporine, dihydroergotamine, ergotamine, fentanyl, pimozide, quinidine, sirolimus, and tacrolimus.
<b>Required Medical Information</b>	Reviewer will verify available claim history to confirm member is not taking simvastatin, lovastatin, cyclosporine, dihydroergotamine, ergotamine, fentanyl, pimozide, quinidine, sirolimus or tacrolimus concurrently with mifepristone.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescriber must be an endocrinologist.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# MIGLUSTAT

---

## Products Affected

- *miglustat*

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	For new starts, documentation of diagnosis for mild to moderate type 1 Gaucher disease. For continuation of therapy or reauthorization: documentation of clinical benefit from use of the drug (i.e. increased platelet count, improvement in anemia, PFT's, improvement in radiographic scans, improved quality of life).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a specialist in treatment of Gaucher's disease
<b>Coverage Duration</b>	New starts will be authorized for 6 months. Cont of therapy or reauth until end of contract year.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# MULTIPLE SCLEROSIS AGENTS

## Products Affected

- BAFIERTAM
- BETASERON SUBCUTANEOUS KIT
- *cladribine (10 tabs)*
- *cladribine (4 tabs)*
- *cladribine (5 tabs)*
- *cladribine (6 tabs)*
- *cladribine (7 tabs)*
- *cladribine (8 tabs)*
- *cladribine (9 tabs)*
- *dimethyl fumarate oral capsule delayed release 120 mg, 240 mg*
- *dimethyl fumarate starter pack oral capsule delayed release therapy pack*
- *fingolimod hcl*
- *glatiramer acetate subcutaneous solution prefilled syringe 20 mg/ml, 40 mg/ml*
- *glatopa subcutaneous solution prefilled syringe 20 mg/ml, 40 mg/ml*
- KESIMPTA
- MAVENCLAD (10 TABS)
- MAVENCLAD (4 TABS)
- MAVENCLAD (5 TABS)
- MAVENCLAD (6 TABS)
- MAVENCLAD (7 TABS)
- MAVENCLAD (8 TABS)
- MAVENCLAD (9 TABS)
- MAYZENT
- MAYZENT STARTER PACK
- PONVORY
- PONVORY STARTER PACK
- REBIF REBIDOSE SUBCUTANEOUS SOLUTION AUTO-INJECTOR
- REBIF REBIDOSE TITRATION PACK SUBCUTANEOUS SOLUTION AUTO-INJECTOR
- REBIF SUBCUTANEOUS SOLUTION PREFILLED SYRINGE
- REBIF TITRATION PACK SUBCUTANEOUS SOLUTION PREFILLED SYRINGE
- TASCENSO ODT
- *teriflunomide*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Specialist for submitted diagnosis.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	If the medication request is for glatiramer, Glatopa, or dimethyl fumarate, the request will be approved. If the member is over 17 years of age and the

<b>PA Criteria</b>	<b>Criteria Details</b>
	request is not for glatiramer, Glatopa, or dimethyl fumarate for multiple sclerosis, the member must have a documented trial of, contraindication to or a medical reason for not using 2 of the following dimethyl fumarate, glatiramer, Glatopa, teriflunomide, or fingolimod. If the request is for fingolimod and the member is 17 years of age or younger, the request will be approved.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# MYFEMBREE

---

## Products Affected

- MYFEMBREE

PA Criteria	Criteria Details
Exclusion Criteria	Patient has history of osteoporosis or hepatic impairment.
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescriber must be an OB, gynecologist or reproductive endocrinologist.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	For new starts for menorrhagia: Trial of, contraindication to, or medical reason for not using an estrogen-progestin contraceptive therapy. For new starts if one of the following drugs has been tried previously, a trial of estrogen-progestin contraceptive therapy is not required: gonadotropin-releasing hormone (GnRH) agonists or tranexamic acid. New starts for endometriosis: Trial of, contraindication to, or medical reason for not using the following concurrently for endometriosis: analgesic pain reliever (e.g. NSAIDs, COX-2 inhibitors) AND either combined estrogen-progestin oral contraceptive, progestin (e.g. medroxyprogesterone acetate, norethindrone), gonadotropin-releasing hormone (GnRH) agonists (e.g. Lupron Depot), OR danazol. For continuation of therapy or reauthorization both of the following are required: 1) Treatment does not exceed the eligible maximum lifetime treatment duration of 2 years, and 2) Documentation has been provided that the member has obtained clinical benefit from medication (e.g. reduced menstrual bleeding from baseline, pain relief).
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Prerequisite Therapy Required</b>	Yes

# NASAL ANTISEIZURE AGENTS

---

## Products Affected

- NAYZILAM
- VALTOCO 10 MG DOSE
- VALTOCO 15 MG DOSE NASAL LIQUID THERAPY PACK 2 X 7.5 MG/0.1ML
- VALTOCO 20 MG DOSE NASAL LIQUID THERAPY PACK 2 X 10 MG/0.1ML
- VALTOCO 5 MG DOSE

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	Nayzilam: 12 years of age or older. Valtoco: 2 years of age or older.
<b>Prescriber Restrictions</b>	Initial therapy only; prescribed by or in consultation with a neurologist.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# NITISINONE

---

## Products Affected

- *nitisinone*
- ORFADIN ORAL SUSPENSION

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescriber must be a geneticist, metabolic specialist, hepatologist, or liver transplant specialist.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	Hereditary Tyrosinemia Type 1 (HT-1): diagnosis was confirmed by genetic testing confirming a mutation of the FAH gene OR elevated levels of succinylacetone.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No
Prerequisite Therapy Required	No

# NON-AMPHETAMINE CENTRAL NERVOUS SYSTEM AGENTS

---

## Products Affected

- *armodafinil*
- *modafinil oral*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No
Prerequisite Therapy Required	No

# NUCALA

## Products Affected

- NUCALA

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Specialist for submitted diagnosis.
<b>Coverage Duration</b>	New starts will be authorized for 6 months. Cont of therapy or reauth until end of contract year.
<b>Other Criteria</b>	New starts for severe asthma: 1) Baseline blood eosinophil count greater than or equal to 150 cells per microliter AND 2) symptoms with equal to or greater than 1 exacerbations in the previous 12 months while on a high-dose inhaled corticosteroid AND 3) the patient has a documented trial of, contraindication to, or medical reason for not using both Dupixent and Fasenra. New starts for eosinophilic granulomatosis with polyangiitis (EGPA): 1) trial of, contraindication to, or medical reason for not using one of the following medications: cyclophosphamide or methotrexate AND 2) the patient has a documented trial of, contraindication to, or medical reason for not using Fasenra. New starts for hypereosinophilic syndrome without an identifiable non-hematologic secondary cause: 1) 2 or more flares within the past 12 months AND 2) trial of, contraindication to, or medical reason for not using oral corticosteroids. New starts for chronic rhinosinusitis with nasal polyps: 1) the patient has a documented trial of, contraindication to, or medical reason for not using both Dupixent and Xolair. Continuation of therapy or re-authorization for all indications: clinical benefit from use of the drug. COPD - initial: (1) documented diagnosis of COPD AND (2) Documentation of blood eosinophil count of greater than or equal to 300 cells/microliter AND (3) Nucala is used as add-on maintenance treatment AND (4) patient has had a trial of, contraindication to, or medical reason for not using at least two of the following: a) long acting beta2-agonist (LABA), b) long acting muscarinic antagonist (LAMA) or c) inhaled

<b>PA Criteria</b>	<b>Criteria Details</b>
	corticosteroid (ICS) AND (5) documentation of COPD exacerbation within the past 12 months. Continuation of therapy: patient has positive clinical response to treatment.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# NUEDEXTA

---

## Products Affected

- NUEDEXTA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	Complete atrioventricular (AV) block without implanted pacemaker, or at high risk of complete AV block. History of heart failure. Concomitant use with MAOIs or use of MAOIs within 14 days. Concomitant use with drugs containing quinidine, quinine, or mefloquine. History of quinine-, mefloquine-, dextromethorphan/quinidine-, or quinidine-induced thrombocytopenia, hepatitis, bone marrow depression, or lupus-like syndrome. Non-Part D indications.
<b>Required Medical Information</b>	Confirmation diagnosis is for Part D indication.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a neurologist or psychiatrist.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# NUPLAZID

---

## Products Affected

- NUPLAZID ORAL CAPSULE
- NUPLAZID ORAL TABLET 10 MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	Patient has a history of dementia-related psychosis.
<b>Required Medical Information</b>	For hallucinations and delusions associated with Parkinson's disease psychosis: documented diagnosis of Parkinson's disease must be made prior to the onset of psychotic symptoms.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# OCTREOTIDE

## Products Affected

- *octreotide acetate injection solution 100 mcg/ml, 1000 mcg/ml, 200 mcg/ml, 50 mcg/ml, 500 mcg/ml*
- *octreotide acetate intramuscular*

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	For new starts for acromegaly: pt meets one of the following (1) inadequate response to surgery and/or radiotherapy OR (2) pt is not an appropriate candidate for surgery and/or radiotherapy OR (3) pt is experiencing negative effects due to tumor size (ex: optic nerve compression). Continuation of therapy or reauthorization: documentation of clinical improvement with therapy.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	Continuation of therapy or reauthorization: documentation of clinical improvement with therapy.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# OFEV

## Products Affected

- *nintedanib esylate*
- OFEV

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescriber must be a pulmonologist, rheumatologist, or lung transplant specialist.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	For a diagnosis of idiopathic pulmonary fibrosis: 1) Documentation of disease as demonstrated on a high resolution CT scan or through lung biopsy and 2) Documented trial of, contraindication to, or medical reason for not using pirfenidone. For a diagnosis of systemic sclerosis-associated interstitial lung disease (SSc-ILD): documented trial of, contraindication to, or medical reason for not using mycophenolate mofetil or cyclophosphamide. For a diagnosis of chronic fibrosing interstitial lung diseases (ILDs) with a progressive phenotype: documentation is provided confirming diagnosis.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# OPSUMIT

---

## Products Affected

- OPSUMIT

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Pulmonary arterial hypertension (PAH) World Health Organization (WHO) Group I - Initial: [Note: documentation required] (1) PAH was confirmed by right heart catheterization AND (2) pretreatment mean pulmonary arterial pressure is greater than 20 mmHg AND (3) pretreatment pulmonary capillary wedge pressure is less than or equal to 15 mmHg AND (4) pretreatment pulmonary vascular resistance is greater than or equal to 3 Wood units. Continuation of therapy: patient has positive clinical response to treatment.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescriber must be a pulmonologist or cardiologist
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	Trial of, contraindication to, or medical reason for not using sildenafil.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# ORAL ANTIPSYCHOTICS

## Products Affected

- CAPLYTA
- COBENFY
- COBENFY STARTER PACK
- FANAPT
- FANAPT TITRATION PACK A
- FANAPT TITRATION PACK B ORAL TABLET
- FANAPT TITRATION PACK C ORAL TABLET
- OPIPZA ORAL FILM 10 MG, 2 MG, 5 MG
- VRAYLAR ORAL CAPSULE

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	Schizophrenia: 1) request is for Caplyta, Cobenfy, Fanapt, Opipza or Vraylar AND 2) trial of, contraindication to, or medical reason for not using two generic antipsychotics. Manic or mixed episodes associated with bipolar I disorder: 1) request is for Fanapt, Opipza or Vraylar AND 2) trial of, contraindication to, or medical reason for not using two generic antipsychotics. Major depressive disorder associated with bipolar I or II disorder: 1) request is for Caplyta AND 2) trial of, contraindication to, or medical reason for not using two generic antipsychotics. Adjunctive treatment of major depressive disorder: 1) request is for Caplyta, Opipza or Vraylar AND 2) trial of, contraindication to, or medical reason for not using two generic antipsychotics AND 3) if the request is for Vraylar: provider attestation that the member is concurrently using an antidepressant. Irritability associated with autistic disorder: 1) request is for Opipza AND 2) trial of, contraindication to, or medical reason for not using two generic antipsychotics. Treatment of Tourette's disorder: 1) request is for Opipza AND 2) trial of, contraindication to, or medical reason for not using two generic antipsychotics. Bipolar I disorder: 1) request is for

<b>PA Criteria</b>	<b>Criteria Details</b>
	Opipza AND 2) trial of, contraindication to, or medical reason for not using two generic antipsychotics. Depressed bipolar I disorder: 1) request is for Vraylar AND 2) trial of, contraindication to, or medical reason for not using two generic antipsychotics.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# ORENCIA

## Products Affected

- ORENCIA CLICKJECT
- ORENCIA INTRAVENOUS
- ORENCIA SUBCUTANEOUS SOLUTION PREFILLED SYRINGE 125 MG/ML, 50 MG/0.4ML, 87.5 MG/0.7ML

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Specialist for submitted diagnosis.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	For pJIA: Either 1) Trial of, medical reason for not using, or contraindication to 2 of the following: Enbrel, an adalimumab product, or Xeljanz or 2) If utilized within the past 120 days, approve for continuation of therapy. For PsA: Either 1) Trial of, medical reason for not using, or contraindication to 2 of the following therapies: an ustekinumab product, Tremfya, Xeljanz, Enbrel, Cosentyx, or an adalimumab product or 2) If utilized within the past 120 days, approve for continuation of therapy. For RA: Either 1) Trial of, medical reason for not using, or contraindication to 2 of the following: Enbrel, an adalimumab product or Xeljanz or 2) If utilized within the past 120 days, approve for continuation of therapy. For acute graft versus host disease: Attestation member is taking in combination with a calcineurin inhibitor and methotrexate.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Prerequisite Therapy Required</b>	Yes

# ORIAHNN

---

## Products Affected

- ORIAHNN

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	Patient has history of osteoporosis or hepatic impairment.
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescriber must be an OB, gynecologist or reproductive endocrinologist.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	For new starts: Trial of, contraindication to, or medical reason for not using an estrogen-progestin contraceptive therapy. For new starts if one of the following drugs has been tried previously, a trial of estrogen-progestin contraceptive therapy is not required: gonadotropin-releasing hormone (GnRH) agonists or tranexamic acid. For continuation of therapy or reauthorization both of the following are required: 1) Treatment does not exceed the eligible maximum lifetime treatment duration of 2 years, and 2) Documentation has been provided that the member has obtained clinical benefit from medication (e.g. reduced menstrual bleeding from baseline).
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# ORILISSA

---

## Products Affected

- ORILISSA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	Patient has osteoporosis or severe hepatic impairment.
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescriber must be an OB or gynecologist.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	Trial of, contraindication to, or medical reason for not using the following concurrently for endometriosis: analgesic pain reliever (e.g. NSAIDs, COX-2 inhibitors) AND either combined estrogen-progestin oral contraceptive, progestin (e.g. medroxyprogesterone acetate, norethindrone), gonadotropin-releasing hormone (GnRH) agonists (e.g. Lupron Depot), OR danazol. For continuation of therapy or reauthorization both of the following are required: 1) Treatment does not exceed the eligible maximum lifetime treatment duration of 2 years for 150mg tablet or 6 months for 200mg tablet, and 2) Documentation has been provided that the member has obtained clinical benefit from the medication.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# ORKAMBI

---

## Products Affected

- ORKAMBI

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	Combination use with Kalydeco, Symdeko, or Trikafta.
<b>Required Medical Information</b>	Documentation of CFTR gene that is responsive to lumacaftor-ivacaftor treatment.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescriber must be a pulmonologist or an expert in the treatment of cystic fibrosis.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# OTEZLA

## Products Affected

- OTEZLA
- OTEZLA XR
- OTEZLA/OTEZLA XR INITIATION PK

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Specialist for submitted diagnosis.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	For moderate to severe psoriasis: Either 1) Trial of, medical reason for not using, or contraindication to 2 of the following therapies: an ustekinumab product, Tremfya, Enbrel, or an adalimumab product or 2) If utilized within the past 120 days, approve for continuation of therapy. For PsA: Either 1) Trial of, medical reason for not using, or contraindication to 2 of the following therapies: an ustekinumab product, Tremfya, Xeljanz, Enbrel, Cosentyx, or an adalimumab product or 2) If utilized within the past 120 days, approve for continuation of therapy. For Behcet's Syndrome or mild psoriasis: Approve.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# OXERVATE

---

## Products Affected

- OXERVATE

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	Treatment duration greater than 16 weeks per affected eye(s).
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by an ophthalmologist or an optometrist.
<b>Coverage Duration</b>	Initial: 8 weeks. Continuation of therapy: additional 8 weeks.
<b>Other Criteria</b>	Initial: confirmed diagnosis. Continuation of therapy: patient has previously received less than or equal to 8 weeks of treatment per affected eye(s) and the patient has a recurrence of neurotrophic keratitis.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# OXYCODONE ER

## Products Affected

- OXYCONTIN ORAL TABLET ER 12 MG, 20 MG, 30 MG, 40 MG, 60 MG, 80 MG  
 HOUR ABUSE-DETERRENT 10 MG, 15 MG

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	Members being treated for active cancer diagnoses, sickle cell diagnoses, those in hospice care, or receiving palliative care will be excluded from the concurrent benzodiazepine and muscle relaxant therapy requirement. For new starts, ALL of the following are required: (1) Member has documented history of receiving an immediate-release opioid, (2) Member has a documented trial of, contraindication to, or medical reason for not using long-acting morphine sulfate, (3) If member is on concurrent benzodiazepines and/or muscle relaxant therapy, the prescriber attests that concurrent therapy is medically necessary, (4) Member is not being treated for substance abuse with buprenorphine-containing products. For continuing therapy, ALL of the following are required: (1) Member's pain has been assessed within the last 6 months, (2) Member has demonstrated clinical improvement in pain and function on current medication regimen, (3) If member is on concurrent benzodiazepines and/or muscle relaxant therapy, the prescriber attests that concurrent therapy is medically necessary, (4) Member is not being treated for substance abuse with buprenorphine-containing products.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# PCSK9 INHIBITORS

---

## Products Affected

- REPATHA
- REPATHA SURECLICK

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	For all diagnoses (besides for risk reduction of major cardiovascular events): documented diagnosis of medically accepted indication. For risk reduction of major cardiovascular (CV) events (CV death, myocardial infarction, stroke, unstable angina requiring hospitalization, or coronary revascularization): 1) Patient has an increased risk of major CV events such as death from CV disease, heart attack, stroke, certain types of chest pain conditions (including but not limited to, unstable angina) requiring hospitalization or certain types of heart surgery and (2) tried or has contraindication to high intensity statin.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# PEGINTERFERON

## Products Affected

- PEGASYS SUBCUTANEOUS SOLUTION 180 MCG/ML
- PEGASYS SUBCUTANEOUS SOLUTION PREFILLED SYRINGE

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	For Hepatitis C: 1) Labs within 3 months of request: liver function tests and detectable HCV RNA viral load. 2) Documentation of genotype, treatment history, and if cirrhotic, documentation of compensated or decompensated cirrhosis. For Hepatitis B: 1) Labs within 3 months of request: ALT/AST, and 2) HBeAg status. For polycythemia vera, approve.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescriber must be a gastroenterologist, hepatologist, infectious disease doctor or transplant specialist.
<b>Coverage Duration</b>	Request will be authorized for 24 to 48 weeks as defined by compendia.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# PENICILLAMINE

---

## Products Affected

- *penicillamine oral tablet*

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Specialist for submitted diagnosis.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	For RA: Trial of, medical reason for not using, or contraindication to 2 of the following: Enbrel, an adalimumab product, or Xeljanz. For other indications, approve.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# PERSERIS

---

## Products Affected

- PERSERIS

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	The member has a documented history of receiving oral risperidone without any clinically significant side effects.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	Trial and failure of, contraindication, or medical reason (e.g. intolerance, hypersensitivity or contraindication) for not using at least two of the following: Abilify Maintena, Abilify Asimtufii, Invega Sustenna, Invega Trinza, Invega Hafyera, or Risperidone Microspheres ER.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# PHENOXYBENZAMINE

---

## Products Affected

- *phenoxybenzamine hcl oral*

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	Trial of, contraindication to, or medical reason for not using doxazosin.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# PIRFENIDONE

---

## Products Affected

- *pirfenidone*

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	For idiopathic pulmonary fibrosis, documentation of all of the following: 1) confirmation of diagnosis on high resolution CT scan or through lung biopsy AND 2) FVC greater than or equal to 50% of the predicted value.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescriber must be a pulmonologist or lung transplant specialist.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# POSACONAZOLE

---

## Products Affected

- *posaconazole oral*

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Documentation of a consultation with an infectious disease specialist, a transplant specialist, or an oncologist.
<b>Coverage Duration</b>	28 days for oropharyngeal candidiasis, end of contract year for other indications
<b>Other Criteria</b>	For treatment of oropharyngeal candidiasis: trial of, contraindication to, or medical reason for not using fluconazole or itraconazole. For prophylaxis of invasive aspergillus infections due to being severely immunocompromised: trial of, contraindication to, or medical reason for not using voriconazole.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# PRETOMANID

---

## Products Affected

- *pretomanid*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	MDR-TB that is not treatment-intolerant or nonresponsive to standard therapy
<b>Required Medical Information</b>	Documentation of use in combination with bedaquiline and linezolid.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an infectious disease specialist.
<b>Coverage Duration</b>	Request will be authorized for 26 weeks.
<b>Other Criteria</b>	Documentation of prior trial of or medical reason for not using first-line TB regimen containing isoniazid and rifampin.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# PREVYMIS

---

## Products Affected

- PREVYMIS ORAL

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescriber must be a hematologist, oncologist, infectious disease, or transplant specialist.
<b>Coverage Duration</b>	Request will be authorized for 6 months.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# PROMACTA

## Products Affected

- *eltrombopag olamine oral packet 12.5 mg, 25 mg*
- *eltrombopag olamine oral tablet 12.5 mg, 25 mg, 50 mg, 75 mg*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	For persistent or chronic immune thrombocytopenia (ITP): Documented baseline platelet count less than 30,000 cells/ microL. For severe aplastic anemia: Documentation of baseline platelet count less than 20,000 cells/microL OR platelet count less than 30,000 cells/microL with bleeding OR reticulocyte count less than 20,000 cells/microL OR absolute neutrophil count less than 500 cells/microL. For thrombocytopenia in patients with Hepatitis C infection: documented baseline platelet count less than 75,000 cells/microL.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Specialist for submitted diagnosis.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	For persistent or chronic immune thrombocytopenia (ITP): Trial of, contraindication to, or medical reason for not using glucocorticosteroids. For severe aplastic anemia: Trial of, contraindication to, or medical reason for not using at least one immunosuppressive agent.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# PYRUKYND

---

## Products Affected

- PYRUKYND
- PYRUKYND TAPER PACK

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	For new starts: 1) documentation of diagnosis and 2) baseline hemoglobin level. For continuation of therapy or reauthorization: documentation of clinical improvement (e.g. reduction in number of blood transfusions, or increase or stabilization in hemoglobin level). If the criteria are not met, may authorize up to 14 days of a Pyrukynd Taper Pack to allow for tapering.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a hematologist.
<b>Coverage Duration</b>	New starts: 6 mo. Cont of therapy or reauth: end of contract yr. Denial: 14 days for dose tapering.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# RADICAVA

## Products Affected

- RADICAVA ORS
- RADICAVA ORS STARTER KIT

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescriber must be a neurologist.
<b>Coverage Duration</b>	New starts: 6 months. Cont. of therapy or reauthorization: until end of contract year.
<b>Other Criteria</b>	For new starts: 1) documentation of ALS functional rating scale (ALSFERS-R) score and 2) documentation that the member has been on riluzole, is beginning therapy as an adjunct to treatment with Radicava, or provider has provided a medical reason why patient is unable to use riluzole. For continuation of therapy or reauthorization: documentation from provider of clinical stabilization in symptoms (e.g. stabilization of ALS functional rating scale (ALSFERS-R) score).
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# RAVICTI

---

## Products Affected

- *glycerol phenylbutyrate*

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Provider is a geneticist, metabolic specialist, gastroenterologist, hepatologist, or liver transplant specialist.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	Trial of, contraindication to, or medical reason for not using sodium phenylbutyrate.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# RECORLEV

---

## Products Affected

- RECORLEV

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Specialist for submitted diagnosis.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	Trial of, contraindication to, or medical reason for not using ketoconazole tablets.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# RELISTOR

## Products Affected

- RELISTOR ORAL
- RELISTOR SUBCUTANEOUS SOLUTION
- RELISTOR SUBCUTANEOUS SOLUTION PREFILLED SYRINGE

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	For opioid induced constipation, chronic non-cancer pain: Patient must have documented trial of or medical reason for not using at least two the following: 1) lubiprostone, OR 2) lactulose OR 3) Movantik. Additionally, patient must have a medical reason for not being able to use oral Relistor in order to receive Relistor injection. For opioid-induced constipation, in patients with advanced illness or pain caused by active cancer requiring opioid dosage escalation for palliative care: Approve.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# REVCOVI

---

## Products Affected

- REVCOVI

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Patient has diagnosis of adenosine deaminase severe combined immune deficiency (ADA-SCID) confirmed by one of the following: 1) documentation of deficiency or absence of adenosine deaminase OR 2) genetic testing revealing mutations in both alleles of the ADA1 gene
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescriber must be an immunologist, geneticist, hematologist, oncologist, or provider who specializes in the treatment of ADA-SCID or related disorders.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# REXULTI

---

## Products Affected

- REXULTI

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	Dementia-related psychosis
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	For schizophrenia: trial of, contraindication to, or medical reason for not using two generic antipsychotics. For major depressive disorder: trial of, contraindication to, or medical reason for not using to two generic antidepressants. For agitation due to dementia: 1) patient has documented diagnosis of Alzheimer's disease AND 2) medication will not be used on an "as needed" basis
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# REZDIFFRA

## Products Affected

- REZDIFFRA

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Patients with decompensated cirrhosis.
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a hepatologist, gastroenterologist, endocrinologist or specialist in the treatment of liver disease.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	Initial therapy: (1) documented diagnosis of noncirrhotic nonalcoholic steatohepatitis (NASH) with moderate to advanced liver fibrosis, (2) documentation of stage F2 to F3 fibrosis as confirmed by a biopsy or a non-invasive test (NIT), (3) the drug is being prescribed at an FDA approved dose according to the member's weight and (4) prescriber attestation to providing lifestyle counseling on nutrition, exercise and avoiding excessive alcohol intake. For reauthorization: (1) the member has shown clinical benefit from the medication (e.g., the resolution of steatohepatitis and no worsening of liver fibrosis, or at least one stage improvement in liver fibrosis and no worsening of steatohepatitis), (2) the member continues to have a fibrosis stage of 3 or less and (3) the drug continues to be prescribed at an FDA approved dose according to the member's weight.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Prerequisite Therapy Required</b>	No

# REZUROCK

---

## Products Affected

- REZUROCK

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescriber must be a hematologist, oncologist, or transplant specialist.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	For new starts: documented trial of, contraindication to, or medical reason for not using at least two lines of systemic immunosuppressive therapy (e.g. corticosteroids, tacrolimus, mycophenolate mofetil, Imbruvica, or Jakafi), one of which must be a systemic corticosteroid. For continuation of therapy or re-authorization: documentation of clinical benefit from use of the drug (i.e. symptom improvement, reduction in corticosteroid dose).
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

# RUFINAMIDE

---

## Products Affected

- *rufinamide oral suspension 40 mg/ml*
- *rufinamide oral tablet*

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	History of familial Short QT syndrome
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescriber must be a neurologist.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	Trial of, contraindication to, or medical reason for not using one alternative generic anticonvulsant for appropriate indications.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# RYKINDO

---

## Products Affected

- RYKINDO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	The member has a documented history of receiving oral risperidone without any clinically significant side effects.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	Trial and failure of, contraindication, or medical reason (e.g. intolerance, hypersensitivity or contraindication) for not using at least two of the following: Abilify Maintena, Abilify Asimtufii, Invega Sustenna, Invega Trinza, or Risperidone Microspheres ER.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# RYLAZE

---

## Products Affected

- RYLAZE

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescriber must be an oncologist, hematologist, or specialist for submitted diagnosis.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# SAPROPTERIN

---

## Products Affected

- *sapropterin dihydrochloride oral packet*
- *sapropterin dihydrochloride oral tablet*

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	For new starts: documentation of elevated baseline phenylalanine levels. Continuation of therapy or reauthorization: prescriber attests the member has improvement in phenylalanine levels from baseline.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	New starts will be authorized for 3 months. Cont of therapy or reauth until end of contract year.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# SECUADO

---

## Products Affected

- SECUADO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	Trial of, contraindication to, or medical reason for not using to one generic antipsychotics.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# SEROSTIM

---

## Products Affected

- SEROSTIM SUBCUTANEOUS SOLUTION RECONSTITUTED 4 MG, 5 MG, 6 MG

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a HIV specialist, gastroenterologist, nutritional support specialist or ID specialist.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	For initial starts for HIV wasting/cachexia: 1) Member must be on anti-retroviral therapy and 2) Alternative causes of wasting have been ruled out (diarrhea, malignancies, inadequate caloric intake, etc)
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

# SIGNIFOR

---

## Products Affected

- SIGNIFOR

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Member is not a candidate for surgery or surgery was not curative.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an endocrinologist.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# SILDENAFIL ORAL

## Products Affected

- *sildenafil citrate oral suspension reconstituted*
- *sildenafil citrate oral tablet 20 mg*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Documentation of concurrent nitrate or Adempas use.
<b>Required Medical Information</b>	Pulmonary Arterial Hypertension: Documentation of pulmonary arterial hypertension (PAH) WHO Group I and PAH Functional Class. Reviewer will verify available patient claim history to confirm patient is not using nitrates or Adempas. Secondary Raynaud's phenomenon- Initial: [Note: documentation required] (1) diagnosis of secondary Raynaud's phenomenon AND (2) diagnosis of primary condition which Raynaud's phenomenon is secondary to (e.g., lupus, scleroderma, rheumatoid arthritis, Sjogren's syndrome, thyroid disease). Continuation of therapy: patient has positive clinical response to treatment.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescriber must be a pulmonologist, cardiologist, or rheumatologist.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	For sildenafil suspension: Documentation of trial of, contraindication to, or medical reason for not using sildenafil tablet.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# SILIQ

---

## Products Affected

- SILIQ

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Specialist for submitted diagnosis.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	For psoriasis: Either 1) Trial of, medical reason for not using, or contraindication to 2 of the following therapies: an ustekinumab product, Tremfya, Enbrel, or an adalimumab product or 2) If utilized within the past 120 days, approve for continuation of therapy.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# SIMPONI

## Products Affected

- SIMPONI SUBCUTANEOUS SOLUTION AUTO-INJECTOR
- SIMPONI SUBCUTANEOUS SOLUTION PREFILLED SYRINGE

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Specialist for submitted diagnosis.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	For ankylosing spondylitis: Either 1) Trial of, medical reason for not using, or contraindication to 2 of the following therapies: Enbrel, an adalimumab product, Cosentyx, or Xeljanz, or 2) If utilized within the past 120 days, approve for continuation of therapy. For PsA: Either 1) Trial of, medical reason for not using, or contraindication to 2 of the following therapies: an ustekinumab product, Cosentyx, Tremfya, Xeljanz, Enbrel, or an adalimumab product or 2) If utilized within the past 120 days, approve for continuation of therapy. For RA: Either 1) Trial of, medical reason for not using, or contraindication to 2 of the following: Enbrel, an adalimumab product, or Xeljanz or 2) If utilized within the past 120 days, approve for continuation of therapy. For adults with UC: Either 1) Trial of, medical reason for not using, or contraindication to 2 of the following: an adalimumab product, an ustekinumab product, Tremfya, or Xeljanz or 2) If utilized within the past 120 days, approve for continuation of therapy. For pediatric patients with UC: approve.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

Formulary ID 26312  
Last Update: 04/21/2026

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Prerequisite Therapy Required</b>	Yes

# SIRTURO

---

## Products Affected

- SIRTURO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Documentation (consistent with pharmacy claims data, OR for new members to the health plan consistent with medical chart history) that the member is currently taking Sirturo as part of a combination regimen with other antimycobacterial drugs to treat MDR-TB.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an infectious disease specialist.
<b>Coverage Duration</b>	Request will be authorized for 24 weeks.
<b>Other Criteria</b>	Documentation of prior trial of or medical reason for not using first-line TB regimen containing isoniazid and rifampin.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# SODIUM OXYBATE

---

## Products Affected

- *sodium oxybate*

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescriber must be a sleep specialist, pulmonologist, or neurologist.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	For somnolence associated with narcolepsy in adults: trial of, contraindication to, or medical reason for not using a CNS stimulant (e.g. methylphenidate, modafinil, armodafinil, etc.). For somnolence associated with narcolepsy in pediatric patients: approve. For cataplexy associated with narcolepsy, approve.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# SODIUM PHENYL BUTYRATE

---

## Products Affected

- *sodium phenylbutyrate oral powder 3 gm/tsp*
- *sodium phenylbutyrate oral tablet*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Provider is a geneticist, metabolic specialist, gastroenterologist, hepatologist, or liver transplant specialist.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# SOFOSBUVIR/VELPATASVIR

---

## Products Affected

- SOFOSBUVIR-VELPATASVIR

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Detectable HCV RNA viral load prior to treatment within 6 months of request. In addition, documentation of treatment history, and if cirrhotic, documentation of compensated or decompensated cirrhosis.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Request will be authorized for 12-24 weeks based on AASLD-IDSA guidelines
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# SOMAVERT

---

## Products Affected

- SOMAVERT

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	For new starts for acromegaly: pt meets one of the following (1) inadequate response to surgery and/or radiotherapy OR (2) pt is not an appropriate candidate for surgery and/or radiotherapy OR (3) pt is experiencing negative effects due to tumor size (ex: optic nerve compression). Continuation of therapy or reauthorization: documentation of clinical improvement with therapy.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an endocrinologist.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# SOTYKTU

---

## Products Affected

- SOTYKTU

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Specialist for submitted diagnosis.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	For moderate to severe psoriasis: Either 1) Trial of, medical reason for not using, or contraindication to 2 of the following therapies: an ustekinumab product, Tremfya, Enbrel, or an adalimumab product or 2) If utilized within the past 120 days, approve for continuation of therapy.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# SUCRAID

---

## Products Affected

- SUCRAID

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	For new starts: documentation of diagnosis of congenital sucrase-isomaltase deficiency. For continuation of therapy or reauthorization: Prescriber attests that member has obtained a clinical benefit (e.g. fewer total stools, greater number of hard and formed stools, fewer watery and soft stools, decrease in breath hydrogen output)
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# SYMDEKO

---

## Products Affected

- SYMDEKO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	Combination use with Kalydeco, Orkambi, or Trikafta.
<b>Required Medical Information</b>	Documentation of CFTR gene that is responsive to tezacaftor-ivacaftor treatment.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescriber must be a pulmonologist or an expert in the treatment of cystic fibrosis.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# SYNAREL

---

## Products Affected

- SYNAREL

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	Trial of, contraindication to, or medical reason for not using the following concurrently for endometriosis: analgesic pain reliever (e.g. NSAIDs, COX-2 inhibitors) AND either combined estrogen-progestin oral contraceptive, progestin (e.g. medroxyprogesterone acetate, norethindrone), OR gonadotropin-releasing hormone (GnRH) agonists (e.g. Lupron Depot)
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# TADALAFIL

---

## Products Affected

- *tadalafil (pah)*
- TADLIQ

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	Documentation of concurrent nitrate or Adempas use.
<b>Required Medical Information</b>	Documentation of pulmonary arterial hypertension (PAH) WHO Group I and PAH Functional Class. Reviewer will verify available patient claim history to confirm patient is not using nitrates or Adempas.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescriber must be a pulmonologist or cardiologist.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	For Tadliq: Documentation of trial of, contraindication to, or medical reason for not using tadalafil tablets.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# TADALAFIL, BPH

---

## Products Affected

- *tadalafil oral tablet 5 mg*

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	Diagnosis of erectile dysfunction
<b>Required Medical Information</b>	Diagnosis of Benign prostatic hyperplasia (BPH) required AND trial of, contraindication to, or medical reason for not using an alpha blocker (e.g. tamsulosin, terazosin).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# TALTZ

## Products Affected

- TALTZ SUBCUTANEOUS SOLUTION AUTO-INJECTOR
- TALTZ SUBCUTANEOUS SOLUTION PREFILLED SYRINGE 20 MG/0.25ML, 40 MG/0.5ML, 80 MG/ML

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Specialist for submitted diagnosis.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	For ankylosing spondylitis: Either 1) Trial of, medical reason for not using, or contraindication to 2 of the following therapies: Enbrel, an adalimumab product, Cosentyx, or Xeljanz, or 2) If utilized within the past 120 days, approve for continuation of therapy. For non-radiographic axial spondyloarthritis: approve. For psoriasis: Either 1) Trial of, medical reason for not using, or contraindication (e.g., safety concerns, not indicated for patient's age) to 2 of the following therapies: an ustekinumab product, Tremfya, Enbrel, or an adalimumab product or 2) If utilized within the past 120 days, approve for continuation of therapy. For PsA: Either 1) Trial of, medical reason for not using, or contraindication to 2 of the following therapies: an ustekinumab product, Cosentyx, Tremfya, Xeljanz, Enbrel, or an adalimumab product or 2) If utilized within the past 120 days, approve for continuation of therapy.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Prerequisite Therapy Required</b>	Yes

# TARPEYO

---

## Products Affected

- TARPEYO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	For new starts: documentation is required that the member has 1) Diagnosis of primary immunoglobulin A nephropathy (IgAN) and 2) at risk of disease progression. Member has proteinuria greater than or equal to 0.5 g/day. For continuation of therapy: documentation that member has been on Tarpeyo for less than 9 months. For reauthorizations: Requests will not be allowed as the safety and efficacy of subsequent courses of Tarpeyo have not been established.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a nephrologist.
<b>Coverage Duration</b>	Request will be authorized for 9 months.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# TASIMELTEON

## Products Affected

- HETLIOZ LQ
- *tasimelteon*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	For new starts of non-24 hour sleep-wake cycle: 1) Member is totally blind with no perception of light, 2) diagnosis of non-24 confirmed by a physiologic circadian phase marker (ex: dim light melatonin onset, assessment of core body temp or measurement of urinary melatonin levels) OR actigraphy with evaluation of sleep logs. For continuation of therapy or reauthorization: documentation of clinical benefit from use of the drug. For night-time sleep disturbances in Smith-Magenis Syndrome (SMS): approve
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Provider is a sleep specialist or neurologist.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# TAVNEOS

## Products Affected

- TAVNEOS

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a rheumatologist or hematologist.
<b>Coverage Duration</b>	New starts will be authorized for 6 months. Cont of therapy or reauth until end of contract year.
<b>Other Criteria</b>	For new starts: 1) Prescriber attests that Tavneos will be prescribed in combination with corticosteroids AND cyclophosphamide unless there is documented trial of, contraindication to, or medical reason for not using these therapies. 2) Documentation of baseline Birmingham Vasculitis Activity Score (BVAS) score 3) Prescriber attestation that the patient will have liver function tests before treatment (ALT, AST, alkaline phosphate, and total bilirubin) and every 4 weeks after start of therapy for the first 6 months of treatment 4) Prescriber attestation that the patient has been screened for and does not have active hepatitis B virus (HBV) infection at baseline. For continuation of therapy or reauthorization: 1) Documentation of remission (BVAS score of 0) OR improvement in BVAS score 2) Prescriber attestation that patient has no abnormality in liver function tests (abnormality: ALT or AST greater than 3 times the upper limit of normal and bilirubin greater than 2 times the upper limit of normal) 3) Prescriber attestation that patient has no active HBV infection.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Prerequisite Therapy Required</b>	Yes

# TEPEZZA

## Products Affected

- TEPEZZA

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Documentation of moderate to severe active thyroid eye disease (TED) as evidenced by one or more of the following: a) eyelid retraction greater than or equal to 2 mm, b) moderate or severe soft tissue involvement, c) exophthalmos greater than or equal to 3 mm above normal for race and sex or d) periodic or constant diplopia OR Documentation of chronic TED with one or more of the following: a) a 3 mm or greater increase in exophthalmos from before diagnosis of TED, b) exophthalmos greater than or equal to 3 mm above normal for race and sex.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescriber must be an ophthalmologist or endocrinologist
<b>Coverage Duration</b>	Request will be authorized for 6 months.
<b>Other Criteria</b>	For active TED: member has had a trial and failure, contraindication to, or medical reason for not using oral or IV glucocorticoids.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# TERIPARATIDE

## Products Affected

- BONSITY
- *teriparatide subcutaneous solution pen-injector 560 mcg/2.24ml*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Documentation showing patient falls into one of the following categories: a bone mineral density (BMD) value consistent with osteoporosis (i.e., T-scores equal to or less than -2.5) or patient has had an osteoporotic fracture or patient has T-scores from -1.5 to -2.5 at the femoral neck or spine, and a 10-year probability of hip fracture greater than or equal to 3% or a 10-year probability of any major osteoporosis-related fracture greater than or equal to 20% based on the United States-adapted FRAX model.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	In addition, the following criteria is also applicable: 1) Trial of, medical reason for not using, or contraindication to an oral bisphosphonate and a denosumab product.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# THIOLA

---

## Products Affected

- *tiopronin oral*

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# TIGECYCLINE

---

## Products Affected

- *tigecycline*

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	(1) Patient must have documented diagnosis of one of the following infections: (a) complicated skin and skin structure infection, (b) complicated intraabdominal infection, (c) community-acquired pneumonia AND (2) Trial of, contraindication to, or medical reason (e.g. intolerance, hypersensitivity or contraindication) for not using at least 2 preferred first-line antibiotics.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an infectious disease specialist.
<b>Coverage Duration</b>	Request will be authorized for 14 days.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# TOBI PODHALER

---

## Products Affected

- TOBI PODHALER

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Patient has documented diagnosis of both: 1) cystic fibrosis AND 2) pseudomonas aeruginosa
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# TOLVAPTAN

---

## Products Affected

- *tolvaptan*
- *tolvaptan (hyponatremia)*

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	Concomitant use with strong CYP3A4 inhibitors (i.e. clarithromycin, ketoconazole, itraconazole, ritonavir, lopinavir-ritonavir, indinavir-ritonavir, indinavir, nelfinavir, saquinavir, nefazodone, conivaptan, and telithromycin).
<b>Required Medical Information</b>	Reviewer will verify available patient claim history to confirm patient is not using a strong CYP3A4 inhibitor (i.e. clarithromycin, ketoconazole, itraconazole, ritonavir, lopinavir-ritonavir, indinavir-ritonavir, indinavir, nelfinavir, saquinavir, nefazodone, conivaptan, and telithromycin).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a cardiologist, endocrinologist, hepatologist, or nephrologist.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# TOPICAL ANTINEOPLASTIC RETINOIDS

---

## Products Affected

- *bexarotene*
- PANRETIN

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a dermatologist, oncologist or specialist for submitted diagnosis
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No
Prerequisite Therapy Required	No

# TOPICAL TESTOSTERONE

## Products Affected

- *testosterone transdermal gel 1.62 %, 12.5 mg/act (1%), 20.25 mg/1.25gm (1.62%), 20.25 mg/act (1.62%), 25 mg/2.5gm (1%), 40.5 mg/2.5gm (1.62%), 50 mg/5gm (1%)*
- *testosterone transdermal solution*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Patient has history of prostate cancer or breast cancer.
<b>Required Medical Information</b>	New starts of topical testosterone therapy for hypogonadism must have both of the following characteristics of hypogonadism: 1) symptoms associated with hypogonadism (e.g. unexplained mild anemia, low libido, decreased energy, etc.) 2) Two separate instances of low serum total or free testosterone taken in the morning, as defined by the lab reference range.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# TRANSDERMAL LIDOCAINE

---

## Products Affected

- *lidocaine external patch 5 %*
- ZTLIDO

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Documented diagnosis of a medically-accepted indication.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	If the request is for the product ZTlido, must provide medical reason for not being able to use generic lidocaine 5% patch
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# TREMFYA

## Products Affected

- TREMFYA ONE-PRESS SUBCUTANEOUS SOLUTION PEN-INJECTOR
- TREMFYA PEN SUBCUTANEOUS SOLUTION AUTO-INJECTOR 100 MG/ML, 200 MG/2ML
- TREMFYA SUBCUTANEOUS SOLUTION PREFILLED SYRINGE 100 MG/ML, 200 MG/2ML
- TREMFYA-CD/UC INDUCTION

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Specialist for submitted diagnosis.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	For UC: Approve. Crohn's Disease (CD) - Initial: Trial of, contraindication to, or medical reason (i.e. patient has a diagnosis of severe Crohn's disease) for not using (i.e. severe Crohns disease), or contraindication to 1 of the following: mercaptopurine, azathioprine, methotrexate, sulfasalazine, or corticosteroid (e.g., prednisone, methylprednisolone). Continuation of therapy: patient has been receiving Tremfya for a minimum of 4 months and has a positive clinical response to therapy.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

# TRIENTINE

---

## Products Affected

- CUVRIOR
- *trientine hcl oral capsule 250 mg*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	If the request is for Cuvrior for new starts, member must have trial of, contraindication to, or medical reason for not using trientine hydrochloride.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

# TRIKAFTA

---

## Products Affected

- TRIKAFTA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	Combination use with Kalydeco, Orkambi, or Symdeko.
<b>Required Medical Information</b>	Documentation of CFTR gene that is responsive to elexacaftor-tezacaftor-ivacaftor treatment.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescriber must be a pulmonologist or an expert in the treatment of cystic fibrosis.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# TYMLOS

---

## Products Affected

- TYMLOS

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Documentation showing patient falls into one of the following categories: a bone mineral density (BMD) value consistent with osteoporosis (i.e., T-scores equal to or less than -2.5) or patient has had an osteoporotic fracture or patient has T-scores from -1.5 to -2.5 at the femoral neck or spine, and a 10-year probability of hip fracture greater than or equal to 3% or a 10-year probability of any major osteoporosis-related fracture greater than or equal to 20% based on the United States-adapted FRAX model.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	The following criteria is also applicable: 1) trial of, contraindication to, or medical reason for not using an oral bisphosphonate and a denosumab product, and 2) therapy does not exceed 2 years.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# TYVASO

---

## Products Affected

- TYVASO DPI MAINTENANCE KIT INHALATION POWDER 112 X 32MCG & 112 X64MCG, 112 X 48MCG & 112 X64MCG, 16 MCG, 32 MCG, 48 MCG, 64 MCG, 80 MCG
- TYVASO DPI TITRATION KIT INHALATION POWDER 16 & 32 & 48 MCG

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescriber must be a pulmonologist or cardiologist.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	For the treatment of pulmonary arterial hypertension (PAH): 1) documentation of PAH WHO Group I classification and PAH Functional Class and 2) trial of, contraindication to, or medical reason for not using a generic phosphodiesterase inhibitor and a generic endothelin receptor antagonist. For the treatment of pulmonary hypertension associated with interstitial lung disease (PH-ILD, WHO Group 3): documentation of PH-ILD and PAH Functional Class.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# UPTRAVI

---

## Products Affected

- UPTRAVI ORAL
- UPTRAVI TITRATION

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Documentation of pulmonary arterial hypertension (PAH) WHO Group I and PAH Functional Class.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescriber must be a pulmonologist or cardiologist.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	Trial of, contraindication to, or medical reason for not using a generic phosphodiesterase inhibitor and a generic endothelin receptor antagonist.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# USTEKINUMAB

## Products Affected

- IMULDOSA SUBCUTANEOUS SOLUTION PREFILLED SYRINGE 45 MG/0.5ML, 90 MG/ML
- SELARSDI INTRAVENOUS
- SELARSDI SUBCUTANEOUS SOLUTION
- SELARSDI SUBCUTANEOUS SOLUTION PREFILLED SYRINGE 45 MG/0.5ML, 90 MG/ML
- STARJEMZA INTRAVENOUS
- STARJEMZA SUBCUTANEOUS SOLUTION
- STARJEMZA SUBCUTANEOUS SOLUTION PREFILLED SYRINGE 45 MG/0.5ML, 90 MG/ML
- STELARA INTRAVENOUS
- STELARA SUBCUTANEOUS SOLUTION 45 MG/0.5ML
- STELARA SUBCUTANEOUS SOLUTION PREFILLED SYRINGE 45 MG/0.5ML, 90 MG/ML
- STEQEYMA INTRAVENOUS
- STEQEYMA SUBCUTANEOUS SOLUTION
- STEQEYMA SUBCUTANEOUS SOLUTION PREFILLED SYRINGE 45 MG/0.5ML, 90 MG/ML
- *ustekinumab subcutaneous solution*
- *ustekinumab subcutaneous solution prefilled syringe 45 mg/0.5ml, 90 mg/ml*
- *ustekinumab-aaaz subcutaneous solution prefilled syringe 45 mg/0.5ml, 90 mg/ml*
- *ustekinumab-aekn subcutaneous solution prefilled syringe 45 mg/0.5ml, 90 mg/ml*
- YESINTEK INTRAVENOUS
- YESINTEK SUBCUTANEOUS SOLUTION
- YESINTEK SUBCUTANEOUS SOLUTION PREFILLED SYRINGE 45 MG/0.5ML, 90 MG/ML

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Specialist for submitted diagnosis.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	N/A
Indications	All Medically-accepted Indications.

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# UZEDY

## Products Affected

- UZEDY SUBCUTANEOUS SUSPENSION PREFILLED SYRINGE 100 MG/0.28ML, 125 MG/0.35ML, 150 MG/0.42ML, 200 MG/0.56ML, 250 MG/0.7ML, 50 MG/0.14ML, 75 MG/0.21ML

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	The member has a documented history of receiving oral risperidone without any clinically significant side effects.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	For schizophrenia: Trial and failure of, contraindication, or medical reason (e.g. intolerance, hypersensitivity or contraindication) for not using at least two of the following: Abilify Maintena, Abilify Asimtufii, Invega Sustenna, Invega Trinza, or Invega Hafyera. For treatment of bipolar I disorder as monotherapy: Trial and failure of, contraindication, or medical reason (e.g. intolerance, hypersensitivity or contraindication) for not using at least one of the following: Abilify Maintena or Abilify Asimtufii. For treatment as adjunctive therapy to lithium or valproate for the maintenance treatment of bipolar I disorder: approve.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# VALCHLOR

---

## Products Affected

- VALCHLOR

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an oncologist or dermatologist.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	Trial of, contraindication to, or medical reason for not being able to use one of the following: a topical corticosteroids or a topical retinoids.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# VEMLIDY

---

## Products Affected

- VEMLIDY

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	For new starts: attestation that member has been tested for HIV infection. If member is HIV-positive, Vemlidy is not used alone.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# VEOZAH

---

## Products Affected

- VEOZAH

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Initial: (1) Documented diagnosis of moderate to severe vasomotor symptoms due to menopause AND (2) Trial of, contraindication to, or medical reason (e.g. intolerance, hypersensitivity or contraindication) for not using a hormonal therapy (e.g., estradiol, oral Premarin, Prempro). Reauthorization: (1) Documentation of positive clinical response to therapy (e.g., decrease in frequency or severity of vasomotor symptoms from baseline)
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# VIGABATRIN

---

## Products Affected

- *vigabatrin*
- VIGAFYDE

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	For infantile spasms or West syndrome, the request will be approved. For diagnosis of refractory complex partial seizures: 1) documentation of diagnosis, and 2) attestation the member is currently receiving another antiepileptic drug, and 3) attestation the member has experienced treatment failure from two alternative antiepileptic agents.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescriber must be a neurologist.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# VIJOICE

---

## Products Affected

- VIJOICE

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	For new starts, all of the following must be included: 1) Documentation of genetic testing confirming diagnosis AND 2) Member has at least one target lesion identified on imaging AND 3) Prescriber attests the patient's condition is severe or life-threatening and necessitates systemic treatment. For continuation of therapy or reauthorization, attestation of a positive clinical response (i.e. reduction in the sum of measurable target lesion volume, absence of progression of non-target lesions, absence of any new lesions, etc.).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a geneticist, dermatologist, vascular surgeon, hematologist/oncologist, or other specialist in the treatment of PIK3CA-Related Overgrowth Spectrum(PROOS).
<b>Coverage Duration</b>	New starts will be authorized for 6 months. Cont of therapy or reauth until end of contract year.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# VMAT-2 INHIBITORS

## Products Affected

- AUSTEDO
- AUSTEDO XR
- AUSTEDO XR PATIENT TITRATION ORAL TABLET EXTENDED RELEASE THERAPY PACK 12 & 18 & 24 & 30 MG
- INGREZZA ORAL CAPSULE
- INGREZZA ORAL CAPSULE SPRINKLE
- INGREZZA ORAL CAPSULE THERAPY PACK
- *tetrabenazine*

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescriber must be a neurologist, clinical geneticist, or psychiatrist.
Coverage Duration	Request will be authorized until the end of the contract year.
Other Criteria	If the request is for tetrabenazine, Ingrezza or Ingrezza Sprinkle, request will be approved. If the request is for Austedo or Austedo XR, the member must have trial of or medical reason for not using tetrabenazine. Reauthorization: Confirmation of improvement in tardive dyskinesia symptoms or chorea associated with Huntington disease symptoms.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

# VORICONAZOLE

---

## Products Affected

- *voriconazole intravenous*

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	Non-Part D indications.
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Request will be authorized for 6 months.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# VOSEVI

---

## Products Affected

- VOSEVI

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Detectable HCV RNA viral load prior to treatment within 6 months of request. In addition, documentation of treatment history, and if cirrhotic, documentation of compensated or decompensated cirrhosis.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Request will be authorized for 12 weeks as per AASLD-IDSA guidance.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# VOWST

---

## Products Affected

- VOWST

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	Treatment of Clostridioides difficile infection (CDI)
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	If all the criteria are met, the request will be approved for 1 month
<b>Other Criteria</b>	Diagnosis of at least 1 recurrent episode of CDI
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# WEGOVY

## Products Affected

- WEGOVY SUBCUTANEOUS SOLUTION AUTO-INJECTOR 0.25 MG/0.5ML, 1.7 MG/0.75ML, 2.4 MG/0.75ML, 0.5 MG/0.5ML, 1

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	The member has an indication of only weight reduction or maintenance for overweight or obesity. The member has concurrent use of any GLP-1 receptor agonist. The member has a personal history of medullary thyroid carcinoma. The member has Multiple Endocrine Neoplasia syndrome type 2.
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	Reduction of the risk of major adverse cardiovascular events in adults with established cardiovascular disease and either obesity or overweight- For new starts: The member has an indication for reducing the risk of adverse cardiovascular events (cardiovascular death, non-fatal myocardial infarction, or non-fatal stroke) in adults with established cardiovascular disease. Documentation demonstrates patient has established cardiovascular disease (e.g., prior myocardial infarction, prior stroke, symptomatic peripheral arterial disease). Documentation is provided that the patient is overweight or obese (defined as a BMI of greater than or equal to 27 kg/m <sup>2</sup> ). Treatment of noncirrhotic metabolic dysfunction-associated steatohepatitis (MASH) - For new starts: The member has a documented diagnosis of MASH with documentation of moderate to advanced liver fibrosis (fibrosis stage 2 or 3) as evidenced by liver biopsy, biomarker testing (AST to platelet ratio index score, fibrosis 4 score, NAFLD fibrosis score, etc.), transient elastography or ultrasonography. For continuation of therapy or reauthorization: Documentation is provided that the patient has a positive clinical response to treatment.

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# WEGOVY TABLETS

## Products Affected

- WEGOVY ORAL

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	The member has an indication of only weight reduction or maintenance for overweight or obesity. The member has concurrent use of any GLP-1 receptor agonist. The member has a personal history of medullary thyroid carcinoma. The member has Multiple Endocrine Neoplasia syndrome type 2.
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	Reduction of the risk of major adverse cardiovascular events in adults with established cardiovascular disease and either obesity or overweight - For new starts: The member has an indication for reducing the risk of adverse cardiovascular events (cardiovascular death, non-fatal myocardial infarction, or non-fatal stroke) in adults with established cardiovascular disease. Documentation demonstrates patient has established cardiovascular disease (i.e., prior myocardial infarction, prior stroke, symptomatic peripheral arterial disease). Documentation is provided that the patient is overweight or obese (defined as a BMI of greater than or equal to 27 kg/m <sup>2</sup> ). The member must have documentation or a medical reason for not being able to use the Wegovy injection. For continuation of therapy or reauthorization: Documentation is provided that the patient has a positive clinical response to treatment.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

Formulary ID 26312

Last Update: 04/21/2026

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Prerequisite Therapy Required</b>	Yes

# WHITE BLOOD CELL STIMULATORS

## Products Affected

- FULPHILA
- FYLNETRA
- LEUKINE INJECTION SOLUTION RECONSTITUTED
- NEULASTA ONPRO SUBCUTANEOUS SOLUTION PREFILLED SYRINGE
- NEULASTA SUBCUTANEOUS SOLUTION 4 MG/0.4ML
- NEULASTA SUBCUTANEOUS SOLUTION PREFILLED SYRINGE
- ZARXIO

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	For new starts for Neulasta and Fylnetra: documentation of trial of, contraindication to, or medical reason for not using Fulphila. Continuation of therapy or re-authorization criteria: diagnosis of chronic neutropenia or a medical reason for continued need for GCSF.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Specialist for submitted diagnosis.
<b>Coverage Duration</b>	New starts will be authorized for 4 months. Cont of therapy or reauth until end of contract year.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# WINREVAIR

## Products Affected

- WINREVAIR

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Pulmonary arterial hypertension (PAH) (World Health Organization [WHO] Group 1)- Initial: [Note: documentation required] (1) PAH was confirmed by right heart catheterization AND (2) pretreatment mean pulmonary arterial pressure is greater than 20 mmHg AND (3) pretreatment pulmonary capillary wedge pressure is less than or equal to 15 mmHg, AND (4) pretreatment pulmonary vascular resistance is greater than or equal to 3 Wood units. Continuation of therapy: patient has positive clinical response to treatment.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a cardiologist or pulmonologist.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	Documented trial and failure of, or contraindication to combination therapy including one PDE-5 inhibitor AND one endothelin receptor antagonist. Documentation of platelet count of greater than 50,000/mm <sup>3</sup> .
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# XDEMVY

---

## Products Affected

- XDEMVY

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# XELJANZ

## Products Affected

- XELJANZ ORAL SOLUTION
- XELJANZ ORAL TABLET
- XELJANZ XR

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Specialist for submitted diagnosis.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	For ankylosing spondylitis: Trial of, medical reason for not using, or contraindication to naproxen and 1 of the following: an adalimumab product, Enbrel, or Cosentyx. For pJIA: Trial of, medical reason for not using, or contraindication to 1 of the following DMARDs: methotrexate or leflunomide and 1 of the following: an adalimumab product or Enbrel. For PsA: Trial of, medical reason for not using, or contraindication to 1 of the following: Enbrel, an adalimumab product, or Cosentyx. For RA: Trial of, medical reason for not using, or contraindication to 1 disease modifying antirheumatic drug (DMARD) (methotrexate, leflunomide, or sulfasalazine) and 1 of the following: Enbrel or an adalimumab product. For UC: Approve. Continuation of therapy: patient has been receiving Xeljanz for a minimum of 4 months and has a positive clinical response.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

Formulary ID 26312  
Last Update: 04/21/2026

# XERMELO

---

## Products Affected

- XERMELO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a gastroenterologist or an oncologist.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	For new starts: 1) Attestation that diarrhea is inadequately controlled by stable dose of SSA therapy for at least three months. For continuation of therapy or reauthorization: 1) documentation of positive clinical response to xermelo and 2) Attestation to continue to be used in combination with SSA.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# XIFAXAN

---

## Products Affected

- XIFAXAN

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	For HE: gastroenterologist or hepatologist. For IBS-D: gastroenterologist.
<b>Coverage Duration</b>	For HE: contract year. For IBSD: 14 days (cannot exceed 3 courses of 14 days each). For TD: 3 days.
<b>Other Criteria</b>	For diagnosis of hepatic encephalopathy (HE): trial of, contraindication to, or medical reason for not using lactulose. For diagnosis of irritable bowel syndrome with diarrhea (IBSD): No more than 3 courses of 14 days each. For travelers diarrhea (TD) caused by noninvasive strains of E. Coli (with no bloody stools or fever): patient must be intolerant to or must have had a trial of at least 3 days of one of the following agents: ciprofloxacin, ofloxacin, levofloxacin or azithromycin.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# XOLAIR

---

## Products Affected

- XOLAIR

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescriber must be a pulmonologist, allergist, immunologist, dermatologist, or otolaryngologist.
Coverage Duration	New starts will be authorized for 6 months. Cont of therapy or reauth until end of contract year.
Other Criteria	New starts for moderate to severe persistent allergic asthma: 1) Evidence of specific allergic sensitivity confirmed by positive skin test (i.e. prick/puncture test) or blood test (i.e. radioallergosorbent test) for a specific IgE or in vitro reactivity to a perennial aeroallergen, AND 2) Pretreatment serum IgE levels greater than 30 IU/mL, AND 3) Symptoms are not adequately controlled with high-dose inhaled corticosteroid (ICS) plus additional controller medication (ie. long-acting B2 agonist) for at least 3 months, or there is a medical reason for not using these drugs. Continuation of therapy or reauthorization criteria for moderate to severe persistent allergic asthma: 1) Reduction in asthma exacerbation resulting in systemic steroid use and/or hospitalization, OR 2) Reduction of rescue inhaler use, OR 3) Documentation of improvement in pulmonary function tests since baseline (prior to initiation of Xolair). New starts for chronic idiopathic urticaria: 1) inadequate symptomatic relief despite trial of two weeks of at least one 2nd generation antihistamine (unless contraindicated), AND 2) disease must be severe enough to warrant short term systemic corticosteroid therapy for management of urticaria. Continuation of therapy or reauthorization criteria for chronic idiopathic urticaria: 1) improvement from baseline of symptoms associated with urticaria within 6 months of Xolair use. New starts for nasal polyps: 1) currently using an intranasal corticosteroid, will be prescribed an intranasal corticosteroid with request, or has a medical reason for not using an intranasal corticosteroid.

<b>PA Criteria</b>	<b>Criteria Details</b>
	Continuation of therapy or reauthorization criteria for nasal polyps: 1) Documentation has been provided that demonstrates a clinical benefit (e.g. improvements in symptom severity, nasal polyp score [NPS], sino-nasal outcome test-22 [SNOT-22], nasal congestion score [NCS]) AND 2) continued use of intranasal corticosteroid, or has a medical reason for not using one. New starts for food allergy: 1) diagnosis of IgE-mediated food allergy 2) Xolair will be used in conjunction with food allergen avoidance. Continuation of therapy or reauthorization criteria for food allergy: 1) Documentation of clinical benefit
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# XOLREMDI

---

## Products Affected

- XOLREMDI

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescriber must be an immunologist or a hematologist
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	For new starts: 1) A documented diagnosis of WHIM (warts, hypogammaglobulinemia, infections and myelokathexis) syndrome confirmed by genotype variant of chemokine receptor 4 (CXCR4) and absolute neutrophil count (ANC) of less than or equal to 400 cells/microliter or white blood cells (WBC) less than or equal to 400 cells/microliter and 2) Documentation of baseline ANC and absolute lymphocyte count (ALC). For renewal 1) Documentation or provider attestation of positive clinical response (i.e. improvement from baseline in ANC, WBC and/or ALC or reduced frequency, duration, or severity of infections, fewer warts, or improved or stabilized clinical signs and/or symptoms of WHIM).
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# XYWAV

---

## Products Affected

- XYWAV

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescriber must be a sleep specialist, pulmonologist or a neurologist.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	For treatment of somnolence associated with narcolepsy, patient must have documentation of either trial of or a medical reason for being unable to use a CNS stimulant (e.g. methylphenidate, modafinil, armodafinil, etc.). For the treatment of cataplexy associated with narcolepsy or idiopathic hypersomnia, approve.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# YORVIPATH

---

## Products Affected

- YORVIPATH

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Diagnosis of acute post-surgical hypoparathyroidism.
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an endocrinologist.
<b>Coverage Duration</b>	New starts will be authorized for 6 months. Cont of therapy or reauth until end of contract year.
<b>Other Criteria</b>	For new starts: 1) Documented diagnosis of chronic hypoparathyroidism AND 2) Provider attestation that patient is currently receiving or has medical reason for not receiving calcium supplementation and active vitamin D treatment AND 3) An albumin-corrected serum calcium level of 7.8 mg/dL or greater. For reauthorization: Documentation of improvement in albumin-corrected serum calcium from baseline.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# YUTREPIA

---

## Products Affected

- YUTREPIA

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a cardiologist or a pulmonologist.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	For the treatment of pulmonary arterial hypertension (PAH): 1) documentation of PAH WHO Group I classification and PAH Functional Class and 2) trial of, contraindication to, or medical reason for not using a generic phosphodiesterase inhibitor and a generic endothelin receptor antagonist. For the treatment of pulmonary hypertension associated with interstitial lung disease (PH-ILD, WHO Group 3): documentation of PHILD and PAH Functional Class.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# ZEPBOUND

## Products Affected

- ZEPBOUND SUBCUTANEOUS SOLUTION AUTO-INJECTOR

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	The member has an indication of only weight reduction or maintenance for overweight or obesity. The member has concurrent use of any GLP-1 receptor agonist. The member has a personal history of medullary thyroid carcinoma. The member has Multiple Endocrine Neoplasia syndrome type 2.
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a sleep disorder specialist, pulmonologist, ENT, or other provider specializing in obstructive sleep apnea.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	For new starts: The member has an indication for moderate to severe obstructive sleep apnea (OSA) in adults with obesity. Documentation of diagnosis of OSA through polysomnography (sleep study) with an apnea-hypopnea index of 15 or more events per hour, or five or more events per hour in the presence of symptoms (e.g., cognitive impairment, fatigue, insomnia, loud snoring) or cardiovascular comorbidities (e.g., hypertension, ischemic heart disease, previous stroke). Documentation is provided that the patient is obese (defined as a BMI of greater than or equal to 30 kg/m <sup>2</sup> ). For continuation of therapy: Documentation of positive response to treatment. Documentation member has achieved and/or maintained a decrease in weight since baseline.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No

Formulary ID 26312  
Last Update: 04/21/2026

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Prerequisite Therapy Required</b>	No

# ZEPOSIA

---

## Products Affected

- ZEPOSIA
- ZEPOSIA 7-DAY STARTER PACK
- ZEPOSIA STARTER KIT ORAL CAPSULE THERAPY PACK 0.23MG & 0.46MG 0.92MG(21)

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Specialist for submitted diagnosis.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	For multiple sclerosis: Trial of, contraindication to, or medical reason for not using two of the following: dalfampridine ER, dimethyl fumarate, fingolimod, glatiramer, glatopa, or teriflunomide. For ulcerative colitis: Either 1) Trial of, medical reason for not using, or contraindication to 1 of the following: an ustekinumab product or an adalimumab product or 2) If utilized within the past 120 days, approve for continuation of therapy.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# ZILBRYSQ

---

## Products Affected

- ZILBRYSQ

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescriber must be a neurologist, rheumatologist, or other appropriate specialist
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	Patient has tried and failed, a medical reason for not using, or has a contraindication to two (2) or more conventional therapies (i.e. pyridostigmine, corticosteroids, or non-steroidal immunosuppressive therapies)
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# ZTALMY

---

## Products Affected

- ZTALMY

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a neurologist.
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year.
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# ZURZUVAE

---

## Products Affected

- ZURZUVAE

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	The member has a documented diagnosis of postpartum depression
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a psychiatrist or obstetrician/gynecologist
<b>Coverage Duration</b>	Request will be authorized until the end of the contract year
<b>Other Criteria</b>	N/A
<b>Indications</b>	All Medically-accepted Indications.
<b>Off-Label Uses</b>	N/A
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

## PART B VERSUS PART D

---

### Products Affected

- *acetylcysteine inhalation solution 10 %, 20 %*
- *acyclovir sodium intravenous solution 50 mg/ml*
- *albuterol sulfate inhalation nebulization solution (2.5 mg/3ml) 0.083%, (5 mg/ml) 0.5%, 0.63 mg/3ml, 1.25 mg/3ml, 2.5 mg/0.5ml*
- *amphotericin b intravenous solution reconstituted 50 mg*
- *amphotericin b liposome intravenous suspension reconstituted 50 mg*
- *aprepitant oral capsule 125 mg, 40 mg, 80 mg*
- *aprepitant oral capsule therapy pack 80 & 125 mg*
- **ASTAGRAF XL ORAL CAPSULE EXTENDED RELEASE 24 HOUR 0.5 MG, 1 MG, 5 MG**
- *azathioprine oral tablet 50 mg*
- *budesonide inhalation suspension 0.25 mg/2ml, 0.5 mg/2ml, 1 mg/2ml*
- *clinisol sf intravenous solution 15 %*
- *cromolyn sodium inhalation nebulization solution 20 mg/2ml*
- *cyclophosphamide oral capsule 25 mg, 50 mg*
- *cyclophosphamide oral tablet 25 mg, 50 mg*
- *cyclosporine modified oral capsule 100 mg, 25 mg, 50 mg*
- *cyclosporine modified oral solution 100 mg/ml*
- *cyclosporine oral capsule 100 mg, 25 mg*
- *dronabinol oral capsule 10 mg, 2.5 mg, 5 mg*
- **EMEND ORAL SUSPENSION RECONSTITUTED 125 MG/5ML**
- **ENGERIX-B INJECTION SUSPENSION 20 MCG/ML**
- **ENGERIX-B INJECTION SUSPENSION PREFILLED SYRINGE 10 MCG/0.5ML, 20 MCG/ML**
- **ENVARUSUS XR ORAL TABLET EXTENDED RELEASE 24 HOUR 0.75 MG, 1 MG, 4 MG**
- *everolimus oral tablet 0.25 mg, 0.5 mg, 0.75 mg, 1 mg*
- *formoterol fumarate inhalation nebulization solution 20 mcg/2ml*
- **GAMMAGARD ERC INJECTION SOLUTION 10 GM/100ML, 5 GM/50ML**
- **GAMMAGARD INJECTION SOLUTION 1 GM/10ML, 10 GM/100ML, 2.5 GM/25ML, 20 GM/200ML, 30 GM/300ML, 5 GM/50ML**
- **GAMMAGARD S/D LESS IGA INTRAVENOUS SOLUTION RECONSTITUTED 10 GM, 5 GM**
- **GAMMAKED INJECTION SOLUTION 1 GM/10ML, 10 GM/100ML, 20 GM/200ML, 5 GM/50ML**
- **GAMMAPLEX INTRAVENOUS SOLUTION 10 GM/100ML, 10 GM/200ML, 20 GM/200ML, 20 GM/400ML, 5 GM/100ML, 5 GM/50ML**
- **GAMUNEX-C INJECTION SOLUTION 1 GM/10ML, 10 GM/100ML, 2.5 GM/25ML, 20 GM/200ML, 40 GM/400ML, 5 GM/50ML**
- *gengraf oral capsule 100 mg, 25 mg*
- *granisetron hcl oral tablet 1 mg*
- **HEPLISAV-B INTRAMUSCULAR SOLUTION PREFILLED SYRINGE 20 MCG/0.5ML**
- **IMOVAX RABIES INTRAMUSCULAR SUSPENSION RECONSTITUTED 2.5 UNIT/ML**
- **INTRALIPID INTRAVENOUS EMULSION 20 %, 30 %**
- *ipratropium bromide inhalation solution 0.02 %*

- *ipratropium-albuterol inhalation solution 0.5-2.5 (3) mg/3ml*
- *levalbuterol hcl inhalation nebulization solution 0.31 mg/3ml, 0.63 mg/3ml, 1.25 mg/3ml*
- *mycophenolate mofetil oral capsule 250 mg*
- *mycophenolate mofetil oral suspension reconstituted 200 mg/ml*
- *mycophenolate mofetil oral tablet 500 mg*
- *mycophenolate sodium oral tablet delayed release 180 mg, 360 mg*
- NUTRILIPID INTRAVENOUS EMULSION 20 %
- OCTAGAM INTRAVENOUS SOLUTION 1 GM/20ML, 10 GM/100ML, 10 GM/200ML, 2 GM/20ML, 2.5 GM/50ML, 20 GM/200ML, 30 GM/300ML, 5 GM/100ML, 5 GM/50ML
- *ondansetron hcl oral solution 4 mg/5ml*
- *ondansetron hcl oral tablet 24 mg, 4 mg, 8 mg*
- *ondansetron oral tablet dispersible 4 mg, 8 mg*
- *pentamidine isethionate inhalation solution reconstituted 300 mg*
- *plenamine intravenous solution 15 %*
- PRIVIGEN INTRAVENOUS SOLUTION 10 GM/100ML, 20 GM/200ML, 40 GM/400ML, 5 GM/50ML
- PROGRAF INTRAVENOUS SOLUTION 5 MG/ML
- PROGRAF ORAL PACKET 0.2 MG, 1 MG
- PULMOZYME INHALATION SOLUTION 2.5 MG/2.5ML
- RABAVERT INTRAMUSCULAR SUSPENSION RECONSTITUTED
- RECOMBIVAX HB INJECTION SUSPENSION 10 MCG/ML, 40 MCG/ML, 5 MCG/0.5ML
- RECOMBIVAX HB INJECTION SUSPENSION PREFILLED SYRINGE 10 MCG/ML, 5 MCG/0.5ML
- *sirolimus oral solution 1 mg/ml*
- *sirolimus oral tablet 0.5 mg, 1 mg, 2 mg*
- *tacrolimus intravenous solution 5 mg/ml*
- *tacrolimus oral capsule 0.5 mg, 1 mg, 5 mg*
- TENIVAC INTRAMUSCULAR INJECTABLE 5-2 LFU (INJECTION)
- TENIVAC INTRAMUSCULAR SUSPENSION 5-2 LF/0.5ML
- *tobramycin inhalation nebulization solution 300 mg/4ml, 300 mg/5ml*

## Details

This drug may be covered under Medicare Part B or D depending upon the circumstances. Information may need to be submitted describing the use and setting of the drug to make the determination.

## Index

### A

abiraterone acetate ..... 11, 13  
 ABIRTEGA ..... 11, 13  
 acetylcysteine inhalation solution 10 %, 20  
 % ..... 236  
 acitretin ..... 1  
 ACTEMRA ACTPEN ..... 4  
 ACTEMRA SUBCUTANEOUS ..... 4  
 acyclovir sodium intravenous solution 50  
 mg/ml ..... 236  
 adalimumab-fkjp (2 pen)..... 5, 6  
 adalimumab-fkjp (2 syringe) subcutaneous  
 prefilled syringe kit 20 mg/0.4ml, 40  
 mg/0.8ml ..... 5, 6  
 ADEMPAS ..... 7  
 AIMOVIG..... 32, 33  
 AKEEGA ..... 11, 13  
 albuterol sulfate inhalation nebulization  
 solution (2.5 mg/3ml) 0.083%, (5 mg/ml)  
 0.5%, 0.63 mg/3ml, 1.25 mg/3ml, 2.5  
 mg/0.5ml ..... 236  
 ALECENSA..... 11, 13  
 ALUNBRIG..... 11, 13  
 ALYFTREK ORAL TABLET 10-50-125  
 MG, 4-20-50 MG ..... 9  
 ambrisentan ..... 10  
 amitriptyline hcl oral ..... 81  
 amoxapine ..... 81  
 amphotericin b intravenous solution  
 reconstituted 50 mg ..... 236  
 amphotericin b liposome intravenous  
 suspension reconstituted 50 mg ..... 236  
 apomorphine hcl subcutaneous ..... 14  
 Aprepitant oral capsule 125 mg, 40 mg, 80  
 mg ..... 236  
 Aprepitant oral capsule therapy pack 80 &  
 125 mg ..... 236  
 AQNEURSA..... 15  
 ARALAST NP INTRAVENOUS  
 SOLUTION RECONSTITUTED 1000  
 MG, 500 MG..... 8  
 ARANESP (ALBUMIN FREE)  
 INJECTION SOLUTION 100 MCG/ML,

200 MCG/ML, 25 MCG/ML, 40  
 MCG/ML, 60 MCG/ML ..... 61  
 ARANESP (ALBUMIN FREE)  
 INJECTION SOLUTION PREFILLED  
 SYRINGE ..... 61  
 ARCALYST ..... 16  
 ARIKAYCE..... 17  
 ARISTADA INITIO ..... 18  
 ARISTADA INTRAMUSCULAR  
 PREFILLED SYRINGE 1064  
 MG/3.9ML, 441 MG/1.6ML, 662  
 MG/2.4ML, 882 MG/3.2ML ..... 18  
 armodafinil ..... 120  
 ASTAGRAF XL ORAL CAPSULE  
 EXTENDED RELEASE 24 HOUR 0.5  
 MG, 1 MG, 5 MG ..... 236  
 AUGTYRO ..... 11, 13  
 AUSTEDO ..... 210  
 AUSTEDO XR ..... 210  
 AUSTEDO XR PATIENT TITRATION  
 ORAL TABLET EXTENDED  
 RELEASE THERAPY PACK 12 & 18 &  
 24 & 30 MG ..... 210  
 AUVELITY ..... 19  
 AVMAPKI FAKZYNJA CO-PACK.. 11, 13  
 AYVAKIT ..... 11, 13  
 azathioprine oral tablet 50 mg..... 236  
**B**  
 bac (butalbital-acetamin-caff)..... 82  
 BAFIERTAM ..... 114, 115  
 BALVERSA ..... 11, 13  
 BENLYSTA SUBCUTANEOUS..... 21, 22  
 benztropine mesylate oral ..... 79, 80  
 BESREMI ..... 23  
 BETASERON SUBCUTANEOUS KIT 114,  
 115  
 bexarotene ..... 11, 13, 193  
 BONSITY ..... 188  
 bosentan oral tablet ..... 24  
 BOSULIF..... 11, 13  
 BRAFTOVI ORAL CAPSULE 75 MG .. 11,  
 13  
 BRINSUPRI..... 25

BRUKINSA ..... 11, 13  
budesonide inhalation suspension 0.25  
mg/2ml, 0.5 mg/2ml, 1 mg/2ml ..... 236  
butalbital-acetaminophen oral tablet 50-325  
mg ..... 82  
butalbital-apap-caff-cod oral capsule 50-  
325-40-30 mg..... 82  
butalbital-apap-caffeine oral capsule 50-  
325-40 mg ..... 82  
butalbital-apap-caffeine oral solution ..... 82  
butalbital-apap-caffeine oral tablet 50-325-  
40 mg ..... 82  
butalbital-asa-caff-codeine..... 82  
butalbital-aspirin-caffeine oral capsule..... 82

**C**  
CABOMETYX ..... 11, 13  
CALQUENCE ORAL TABLET ..... 11, 13  
CAMZYOS ..... 26, 27  
CAPLYTA ..... 128, 129  
CAPRELSA ..... 11, 13  
CARDAMYST ..... 28  
carglumic acid oral tablet soluble ..... 29  
carisoprodol oral ..... 84  
casprofungin acetate..... 30  
CAYSTON..... 20  
CERDELGA ..... 31  
chlorzoxazone oral tablet 500 mg ..... 84  
CHOLBAM..... 34  
CIBINQO ..... 35  
CIMZIA (1 SYRINGE) ..... 36, 37  
CIMZIA (2 SYRINGE) ..... 36, 37  
CIMZIA SUBCUTANEOUS KIT 2 X 200  
MG ..... 36, 37  
CIMZIA-STARTER ..... 36, 37  
CINRYZE ..... 78  
cladribine (10 tabs) ..... 114, 115  
cladribine (4 tabs) ..... 114, 115  
cladribine (5 tabs) ..... 114, 115  
cladribine (6 tabs) ..... 114, 115  
cladribine (7 tabs) ..... 114, 115  
cladribine (8 tabs) ..... 114, 115  
cladribine (9 tabs) ..... 114, 115  
clinisol sf intravenous solution 15 % ..... 236  
clomipramine hcl oral ..... 81  
COBENFY ..... 128, 129  
COBENFY STARTER PACK ..... 128, 129

COMETRIQ (100 MG DAILY DOSE)  
ORAL KIT 80 & 20 MG ..... 11, 13  
COMETRIQ (140 MG DAILY DOSE)  
ORAL KIT 3 X 20 MG & 80 MG .. 11, 13  
COMETRIQ (60 MG DAILY DOSE) 11, 13  
COPIKTRA..... 11, 13  
CORLANOR ORAL SOLUTION..... 38  
CORTROPHIN ..... 39  
CORTROPHIN GEL ..... 39  
COSENTYX (300 MG DOSE)..... 40  
COSENTYX INTRAVENOUS..... 40  
COSENTYX SENSOREADY (300 MG). 40  
COSENTYX SENSOREADY PEN ..... 40  
COSENTYX SUBCUTANEOUS  
SOLUTION PREFILLED SYRINGE 150  
MG/ML, 75 MG/0.5ML ..... 40  
COSENTYX UNOREADY ..... 40  
COTELLIC ..... 11, 13  
CRESEMBA ORAL ..... 41  
cromolyn sodium inhalation nebulization  
solution 20 mg/2ml ..... 236  
CUVRIOR..... 197  
cyclobenzaprine hcl oral tablet 10 mg, 5 mg  
..... 84  
cyclophosphamide oral capsule 25 mg, 50  
mg ..... 236  
cyclophosphamide oral tablet 25 mg, 50 mg  
..... 236  
cyclosporine modified oral capsule 100 mg,  
25 mg, 50 mg ..... 236  
cyclosporine modified oral solution 100  
mg/ml ..... 236  
cyclosporine oral capsule 100 mg, 25 mg 236  
cyproheptadine hcl oral..... 79, 80  
CYSTAGON..... 42  
CYSTARAN ..... 43

**D**  
dalfampridine er ..... 44  
DANZITEN ..... 11, 13  
dasatinib ..... 11, 13  
DAURISMO ..... 11, 13  
deferasirox..... 45  
deferasirox granules ..... 45  
deferiprone ..... 46  
DIACOMIT..... 48  
dihydroergotamine mesylate nasal..... 49

dimethyl fumarate oral capsule delayed  
 release 120 mg, 240 mg ..... 114, 115  
 dimethyl fumarate starter pack oral capsule  
 delayed release therapy pack ..... 114, 115  
 diphenoxylate-atropine oral liquid..... 79, 80  
 diphenoxylate-atropine oral tablet 2.5-0.025  
 mg ..... 79, 80  
 dipyridamole oral ..... 79, 80  
 DOPTELET ..... 50  
 DOPTELET SPRINKLE ..... 50  
 doxepin hcl external..... 51  
 dronabinol oral capsule 10 mg, 2.5 mg, 5 mg  
 ..... 236  
 DUPIXENT SUBCUTANEOUS  
 SOLUTION AUTO-INJECTOR .... 52, 53  
 DUPIXENT SUBCUTANEOUS  
 SOLUTION PREFILLED SYRINGE 200  
 MG/1.14ML, 300 MG/2ML ..... 52, 53  
**E**  
 EGRIFTA SV ..... 54  
 EGRIFTA WR ..... 54  
 ELIGARD ..... 75  
 eltrombopag olamine oral packet 12.5 mg,  
 25 mg ..... 148  
 eltrombopag olamine oral tablet 12.5 mg, 25  
 mg, 50 mg, 75 mg ..... 148  
 EMEND ORAL SUSPENSION  
 RECONSTITUTED 125 MG/5ML .... 236  
 EMGALITY ..... 32, 33  
 EMGALITY (300 MG DOSE) ..... 32, 33  
 EMSAM ..... 55  
 ENBREL MINI ..... 56  
 ENBREL SUBCUTANEOUS SOLUTION  
 25 MG/0.5ML ..... 56  
 ENBREL SUBCUTANEOUS SOLUTION  
 PREFILLED SYRINGE ..... 56  
 ENBREL SURECLICK SUBCUTANEOUS  
 SOLUTION AUTO-INJECTOR ..... 56  
 ENGERIX-B INJECTION SUSPENSION  
 20 MCG/ML ..... 236  
 ENGERIX-B INJECTION SUSPENSION  
 PREFILLED SYRINGE 10 MCG/0.5ML,  
 20 MCG/ML ..... 236  
 ENSACOVE ..... 11, 13  
 ENTYVIO PEN ..... 58

ENVARUSUS XR ORAL TABLET  
 EXTENDED RELEASE 24 HOUR 0.75  
 MG, 1 MG, 4 MG ..... 236  
 EPIDIOLEX..... 59  
 EPOGEN INJECTION SOLUTION 10000  
 UNIT/ML, 2000 UNIT/ML, 20000  
 UNIT/ML, 3000 UNIT/ML, 4000  
 UNIT/ML ..... 61  
 ergotamine-caffeine ..... 79, 80  
 ERIVEDGE..... 11, 13  
 ERLEADA ..... 11, 13  
 erlotinib hcl ..... 11, 13  
 ERZOFRI INTRAMUSCULAR  
 SUSPENSION PREFILLED SYRINGE  
 117 MG/0.75ML, 156 MG/ML, 234  
 MG/1.5ML, 351 MG/2.25ML, 39  
 MG/0.25ML, 78 MG/0.5ML ..... 62  
 eszopiclone..... 85  
 EUCRISA ..... 63  
 EULEXIN ..... 11, 13  
 everolimus oral tablet 0.25 mg, 0.5 mg, 0.75  
 mg, 1 mg ..... 236  
 everolimus oral tablet 10 mg, 2.5 mg, 5 mg,  
 7.5 mg ..... 11, 13  
 everolimus oral tablet soluble ..... 11, 13  
 EVRYSDI ..... 64  
**F**  
 FABHALTA ..... 65  
 FANAPT ..... 128, 129  
 FANAPT TITRATION PACK A ... 128, 129  
 FANAPT TITRATION PACK B ORAL  
 TABLET ..... 128, 129  
 FANAPT TITRATION PACK C ORAL  
 TABLET ..... 128, 129  
 FASENRA ..... 66, 67  
 FASENRA PEN ..... 66, 67  
 FILSPARI ..... 68  
 fingolimod hcl ..... 114, 115  
 FINTEPLA ..... 69  
 FIRDAPSE ..... 70  
 FIRMAGON (240 MG DOSE) ..... 75  
 FIRMAGON SUBCUTANEOUS  
 SOLUTION RECONSTITUTED 80 MG  
 ..... 75  
 flucytosine oral ..... 71

formoterol fumarate inhalation nebulization solution 20 mcg/2ml ..... 236

FOTIVDA ..... 11, 13

FRUZAQLA ..... 11, 13

FULPHILA ..... 218

FYLNETHA ..... 218

**G**

GALAFOLD ..... 72

GAMMAGARD ERC INJECTION SOLUTION 10 GM/100ML, 5 GM/50ML ..... 236

GAMMAGARD INJECTION SOLUTION 1 GM/10ML, 10 GM/100ML, 2.5 GM/25ML, 20 GM/200ML, 30 GM/300ML, 5 GM/50ML ..... 236

GAMMAGARD S/D LESS IGA INTRAVENOUS SOLUTION RECONSTITUTED 10 GM, 5 GM .... 236

GAMMAKED INJECTION SOLUTION 1 GM/10ML, 10 GM/100ML, 20 GM/200ML, 5 GM/50ML ..... 236

GAMMAPLEX INTRAVENOUS SOLUTION 10 GM/100ML, 10 GM/200ML, 20 GM/200ML, 20 GM/400ML, 5 GM/100ML, 5 GM/50ML ..... 236

GAMUNEX-C INJECTION SOLUTION 1 GM/10ML, 10 GM/100ML, 2.5 GM/25ML, 20 GM/200ML, 40 GM/400ML, 5 GM/50ML ..... 236

GATTEX ..... 73

GAVRETO ..... 11, 13

gefitinib ..... 11, 13

gengraf oral capsule 100 mg, 25 mg ..... 236

GENOTROPIN MINIQUICK SUBCUTANEOUS PREFILLED SYRINGE ..... 77

GENOTROPIN SUBCUTANEOUS CARTRIDGE ..... 77

GILOTRIF ..... 11, 13

GLASSIA ..... 8

glatiramer acetate subcutaneous solution prefilled syringe 20 mg/ml, 40 mg/ml 114, 115

glatopa subcutaneous solution prefilled syringe 20 mg/ml, 40 mg/ml ..... 114, 115

glyburide micronized oral tablet 3 mg, 6 mg ..... 79, 80

glyburide oral tablet 1.25 mg, 2.5 mg, 5 mg ..... 79, 80

glyburide-metformin oral tablet 1.25-250 mg ..... 79, 80

glyburide-metformin oral tablet 2.5-500 mg, 5-500 mg ..... 79, 80

glycerol phenylbutyrate ..... 151

GOCOVRI ..... 76

GOMEKLI ..... 11, 13

granisetron hcl oral tablet 1 mg ..... 236

guanfacine hcl er oral tablet extended release 24 hour 1 mg, 2 mg, 3 mg, 4 mg ..... 79, 80

guanfacine hcl oral ..... 79, 80

**H**

HAEGARDA ..... 78

HEPLISAV-B INTRAMUSCULAR SOLUTION PREFILLED SYRINGE 20 MCG/0.5ML ..... 236

HERNEXEOS ..... 11, 13

HETLIOZ LQ ..... 184

hydroxyzine hcl oral syrup 10 mg/5ml 79, 80

hydroxyzine hcl oral tablet ..... 79, 80

hydroxyzine pamoate oral ..... 79, 80

HYFTOR ..... 86

HYRNUO ..... 11, 13

**I**

IBRANCE ..... 11, 13

IBTROZI ..... 11, 13

icatibant acetate subcutaneous solution prefilled syringe ..... 87

ICLUSIG ..... 11, 13

IDHIFA ..... 11, 13

ILARIS SUBCUTANEOUS SOLUTION 88

ILUMYA ..... 89

imatinib mesylate oral ..... 11, 13

IMBRUVICA ORAL CAPSULE ..... 90

IMBRUVICA ORAL SUSPENSION ..... 90

IMBRUVICA ORAL TABLET 140 MG, 280 MG, 420 MG ..... 90

IMKELDI ..... 11, 13

IMOVAX RABIES INTRAMUSCULAR SUSPENSION RECONSTITUTED 2.5 UNIT/ML ..... 236

IMPAVIDO.....	91
IMULDOSA SUBCUTANEOUS SOLUTION PREFILLED SYRINGE 45 MG/0.5ML, 90 MG/ML .....	202, 203
INCRELEX.....	92
indomethacin oral capsule 25 mg, 50 mg	79, 80
INGREZZA ORAL CAPSULE .....	210
INGREZZA ORAL CAPSULE SPRINKLE .....	210
INGREZZA ORAL CAPSULE THERAPY PACK.....	210
INLURIYO .....	11, 13
INLYTA.....	11, 13
INQOVI .....	11, 13
INREBIC.....	11, 13
INTRALIPID INTRAVENOUS EMULSION 20 %, 30 % .....	236
ipratropium bromide inhalation solution 0.02 % .....	236
ipratropium-albuterol inhalation solution 0.5-2.5 (3) mg/3ml .....	237
ITOVEBI.....	11, 13
ivabradine hcl.....	38
IWILFIN .....	11, 13
<b>J</b>	
JAKAFI.....	93
JAYPIRCA .....	11, 13
JYLAMVO .....	109
<b>K</b>	
KALYDECO.....	94
KERENDIA .....	95, 96
KESIMPTA.....	114, 115
ketorolac tromethamine oral .....	79, 80
KEVZARA .....	97
KINERET SUBCUTANEOUS SOLUTION PREFILLED SYRINGE .....	98
KISQALI (200 MG DOSE) .....	11, 13
KISQALI (400 MG DOSE) .....	11, 13
KISQALI (600 MG DOSE) .....	11, 13
KISQALI FEMARA (400 MG DOSE) ...	11, 13
KISQALI FEMARA (600 MG DOSE) ...	11, 13
KOMZIFTI .....	11, 13
KOSELUGO .....	11, 13

KRAZATI.....	11, 13
KYMBEE.....	47
<b>L</b>	
lapatinib ditosylate .....	11, 13
LAZCLUZE .....	11, 13
lenalidomide.....	11, 13
LENVIMA (10 MG DAILY DOSE) ..	11, 13
LENVIMA (12 MG DAILY DOSE) ..	11, 13
LENVIMA (14 MG DAILY DOSE) ..	11, 13
LENVIMA (18 MG DAILY DOSE) ..	11, 13
LENVIMA (20 MG DAILY DOSE) ..	11, 13
LENVIMA (24 MG DAILY DOSE) ..	11, 13
LENVIMA (4 MG DAILY DOSE) ....	11, 13
LENVIMA (8 MG DAILY DOSE) ....	11, 13
LEQEMBI IQLIK.....	99
LEQSELVI .....	100
LEUKERAN .....	11, 13
LEUKINE INJECTION SOLUTION RECONSTITUTED .....	218
leuprolide acetate (3 month) .....	75
leuprolide acetate injection .....	75
levalbuterol hcl inhalation nebulization solution 0.31 mg/3ml, 0.63 mg/3ml, 1.25 mg/3ml .....	237
l-glutamine oral packet .....	57
lidocaine external patch 5 % .....	195
liraglutide .....	74
LITFULO.....	101
LIVMARLI.....	102
LIVTENCITY.....	103
LODOCO.....	104
lofexidine hcl .....	105
LONSURF .....	11, 13
LORBRENA.....	11, 13
LUMAKRAS .....	11, 13
LUPKYNIS.....	106
LUPRON DEPOT (1-MONTH) .....	75
LUPRON DEPOT (3-MONTH) .....	75
LUPRON DEPOT (4-MONTH) .....	75
LUPRON DEPOT (6-MONTH) .....	75
LUTRATE DEPOT .....	75
LYBALVI.....	107
LYNPARZA ORAL TABLET .....	11, 13
LYTGObI (12 MG DAILY DOSE) ...	11, 13
LYTGObI (16 MG DAILY DOSE) ...	11, 13
LYTGObI (20 MG DAILY DOSE) ...	11, 13

**M**

MAVENCLAD (10 TABS) ..... 114, 115  
MAVENCLAD (4 TABS) ..... 114, 115  
MAVENCLAD (5 TABS) ..... 114, 115  
MAVENCLAD (6 TABS) ..... 114, 115  
MAVENCLAD (7 TABS) ..... 114, 115  
MAVENCLAD (8 TABS) ..... 114, 115  
MAVENCLAD (9 TABS) ..... 114, 115  
MAVYRET ..... 108  
MAYZENT ..... 114, 115  
MAYZENT STARTER PACK ..... 114, 115  
megestrol acetate oral suspension 40 mg/ml,  
400 mg/10ml, 625 mg/5ml ..... 83  
megestrol acetate oral tablet ..... 83  
MEKINIST ..... 11, 13  
MEKTOVI ..... 11, 13  
mercaptopurine oral suspension ..... 12, 13  
metaxalone oral tablet 800 mg ..... 84  
methocarbamol oral tablet 500 mg, 750 mg  
..... 84  
methyltestosterone oral ..... 110  
metyrosine ..... 111  
mifepristone oral tablet 300 mg ..... 112  
miglustat ..... 113  
modafinil oral ..... 120  
MODEYSO ..... 12, 13  
MOUNJARO SUBCUTANEOUS  
SOLUTION AUTO-INJECTOR ..... 74  
mycophenolate mofetil oral capsule 250 mg  
..... 237  
mycophenolate mofetil oral suspension  
reconstituted 200 mg/ml ..... 237  
mycophenolate mofetil oral tablet 500 mg  
..... 237  
mycophenolate sodium oral tablet delayed  
release 180 mg, 360 mg ..... 237  
MYFEMBREE ..... 116, 117

**N**  
NAYZILAM ..... 118  
NERLYNX ..... 12, 13  
NEULASTA ONPRO SUBCUTANEOUS  
SOLUTION PREFILLED SYRINGE 218  
NEULASTA SUBCUTANEOUS  
SOLUTION 4 MG/0.4ML ..... 218  
NEULASTA SUBCUTANEOUS  
SOLUTION PREFILLED SYRINGE 218

NEXLETOL ..... 2, 3  
NEXLIZET ..... 2, 3  
NGENLA ..... 77  
nifedipine oral ..... 79, 80  
nilotinib d-tartrate ..... 12, 13  
nilotinib hcl ..... 12, 13  
nilutamide ..... 12, 13  
NINLARO ..... 12, 13  
nintedanib esylate ..... 126  
nitisinone ..... 119  
nortriptyline hcl oral ..... 81  
NUBEQA ..... 12, 13  
NUCALA ..... 121, 122  
NUEDEXTA ..... 123  
NUPLAZID ORAL CAPSULE ..... 124  
NUPLAZID ORAL TABLET 10 MG .... 124  
NURTEC ..... 32, 33  
NUTRILIPID INTRAVENOUS  
EMULSION 20 % ..... 237

**O**  
OCTAGAM INTRAVENOUS SOLUTION  
1 GM/20ML, 10 GM/100ML, 10  
GM/200ML, 2 GM/20ML, 2.5  
GM/50ML, 20 GM/200ML, 30  
GM/300ML, 5 GM/100ML, 5 GM/50ML  
..... 237  
octreotide acetate injection solution 100  
mcg/ml, 1000 mcg/ml, 200 mcg/ml, 50  
mcg/ml, 500 mcg/ml ..... 125  
octreotide acetate intramuscular ..... 125  
ODOMZO ..... 12, 13  
OFEV ..... 126  
OGSIVEO ORAL TABLET 100 MG, 150  
MG ..... 12, 13  
OJEMDA ..... 12, 13  
OJJAARA ..... 12, 13  
OMNITROPE SUBCUTANEOUS  
SOLUTION CARTRIDGE ..... 77  
OMNITROPE SUBCUTANEOUS  
SOLUTION RECONSTITUTED ..... 77  
ondansetron hcl oral solution 4 mg/5ml.. 237  
ondansetron hcl oral tablet 24 mg, 4 mg, 8  
mg ..... 237  
ondansetron oral tablet dispersible 4 mg, 8  
mg ..... 237  
ONUREG ..... 12, 13

OPIPZA ORAL FILM 10 MG, 2 MG, 5 MG ..... 128, 129  
 OPSUMIT ..... 127  
 ORENCIA CLICKJECT..... 130, 131  
 ORENCIA INTRAVENOUS ..... 130, 131  
 ORENCIA SUBCUTANEOUS SOLUTION  
 PREFILLED SYRINGE 125 MG/ML, 50  
 MG/0.4ML, 87.5 MG/0.7ML ..... 130, 131  
 ORFADIN ORAL SUSPENSION..... 119  
 ORGOVYX..... 12, 13  
 ORIAHNN ..... 132  
 ORLISSA..... 133  
 ORKAMBI..... 134  
 ORLADEYO..... 78  
 orphenadrine citrate er ..... 84  
 ORSERDU ..... 12, 13  
 OTEZLA ..... 135  
 OTEZLA XR ..... 135  
 OTEZLA/OTEZLA XR INITIATION PK  
 ..... 135  
 OXERVATE ..... 136  
 OXYCONTIN ORAL TABLET ER 12  
 HOUR ABUSE-DETERRENT 10 MG,  
 15 MG, 20 MG, 30 MG, 40 MG, 60 MG,  
 80 MG ..... 137, 138  
 OZEMPIC (0.25 OR 0.5 MG/DOSE)  
 SUBCUTANEOUS SOLUTION PEN-  
 INJECTOR 2 MG/3ML ..... 74  
 OZEMPIC (1 MG/DOSE)  
 SUBCUTANEOUS SOLUTION PEN-  
 INJECTOR 4 MG/3ML ..... 74  
 OZEMPIC (2 MG/DOSE) ..... 74  
**P**  
 PANRETIN..... 193  
 pazopanib hcl ..... 12, 13  
 PEGASYS SUBCUTANEOUS SOLUTION  
 180 MCG/ML ..... 140  
 PEGASYS SUBCUTANEOUS SOLUTION  
 PREFILLED SYRINGE ..... 140  
 PEMAZYRE ..... 12, 13  
 penicillamine oral tablet..... 141  
 pentamidine isethionate inhalation solution  
 reconstituted 300 mg..... 237  
 perphenazine-amitriptyline ..... 81  
 PERSERIS ..... 142  
 phenobarbital oral elixir ..... 81

phenobarbital oral tablet ..... 81  
 phenoxybenzamine hcl oral ..... 143  
 PIQRAY (200 MG DAILY DOSE).... 12, 13  
 PIQRAY (250 MG DAILY DOSE).... 12, 13  
 PIQRAY (300 MG DAILY DOSE).... 12, 13  
 pirfenidone ..... 144  
 plenamine intravenous solution 15 %..... 237  
 pomalidomide ..... 12, 13  
 PONVORY ..... 114, 115  
 PONVORY STARTER PACK..... 114, 115  
 posaconazole oral..... 145  
 pretomanid ..... 146  
 PREVYMIS ORAL ..... 147  
 PRIVIGEN INTRAVENOUS SOLUTION  
 10 GM/100ML, 20 GM/200ML, 40  
 GM/400ML, 5 GM/50ML ..... 237  
 PROCRIT..... 61  
 PROGRAF INTRAVENOUS SOLUTION 5  
 MG/ML ..... 237  
 PROGRAF ORAL PACKET 0.2 MG, 1 MG  
 ..... 237  
 PROLASTIN-C INTRAVENOUS  
 SOLUTION..... 8  
 promethazine hcl oral solution 6.25 mg/5ml  
 ..... 79, 80  
 promethazine hcl oral tablet..... 79, 80  
 promethazine hcl rectal suppository 12.5  
 mg, 25 mg ..... 79, 80  
 promethazine-phenylephrine..... 79, 80  
 promethegan rectal suppository 50 mg 79, 80  
 PULMOZYME INHALATION  
 SOLUTION 2.5 MG/2.5ML ..... 237  
 PYRUKYND ..... 149  
 PYRUKYND TAPER PACK..... 149  
**Q**  
 QINLOCK..... 12, 13  
 QULIPTA ..... 32, 33  
**R**  
 RABAVERT INTRAMUSCULAR  
 SUSPENSION RECONSTITUTED... 237  
 RADICAVA ORS..... 150  
 RADICAVA ORS STARTER KIT ..... 150  
 REBIF REBIDOSE SUBCUTANEOUS  
 SOLUTION AUTO-INJECTOR 114, 115

REBIF REBIDOSE TITRATION PACK SUBCUTANEOUS SOLUTION AUTO- INJECTOR.....	114, 115	sapropterin dihydrochloride oral tablet... 162
REBIF SUBCUTANEOUS SOLUTION PREFILLED SYRINGE.....	114, 115	SCSEMBLIX .....
REBIF TITRATION PACK SUBCUTANEOUS SOLUTION PREFILLED SYRINGE .....	114, 115	SECUADO.....
RECOMBIVAX HB INJECTION SUSPENSION 10 MCG/ML, 40 MCG/ML, 5 MCG/0.5ML .....	237	SELARSDI INTRAVENOUS .....
RECOMBIVAX HB INJECTION SUSPENSION PREFILLED SYRINGE 10 MCG/ML, 5 MCG/0.5ML .....	237	SELARSDI SUBCUTANEOUS SOLUTION.....
RECORLEV .....	152	SELARSDI SUBCUTANEOUS SOLUTION PREFILLED SYRINGE 45 MG/0.5ML, 90 MG/ML .....
RELISTOR ORAL.....	153	202, 203
RELISTOR SUBCUTANEOUS SOLUTION.....	153	SEROSTIM SUBCUTANEOUS SOLUTION RECONSTITUTED 4 MG, 5 MG, 6 MG.....
RELISTOR SUBCUTANEOUS SOLUTION PREFILLED SYRINGE	153	164
REPATHA .....	139	SIGNIFOR .....
REPATHA SURECLICK.....	139	165
RETACRIT INJECTION SOLUTION 10000 UNIT/ML, 10000 UNIT/ML(1ML), 2000 UNIT/ML, 20000 UNIT/ML, 3000 UNIT/ML, 4000 UNIT/ML, 40000 UNIT/ML .....	61	sildenafil citrate oral suspension reconstituted.....
RETEVMO ORAL TABLET.....	12, 13	166
REVCovi.....	154	sildenafil citrate oral tablet 20 mg .....
REVLIMID.....	12, 13	166
REVUFORJ .....	12, 13	SILIQ .....
REXULTI .....	155	167
REZDIFFRA.....	156, 157	SIMLANDI (1 PEN) SUBCUTANEOUS AUTO-INJECTOR KIT 40 MG/0.4ML, 80 MG/0.8ML .....
REZLIDHIA .....	12, 13	5, 6
REZUROCK.....	158	SIMLANDI (1 SYRINGE).....
ROMVIMZA .....	12, 13	5, 6
ROZLYTREK.....	12, 13	SIMLANDI (2 PEN).....
RUBRACA .....	12, 13	5, 6
rufinamide oral suspension 40 mg/ml.....	159	SIMLANDI (2 SYRINGE) SUBCUTANEOUS PREFILLED SYRINGE KIT 20 MG/0.2ML, 40 MG/0.4ML .....
rufinamide oral tablet.....	159	5, 6
RYBELSUS .....	74	SIMPONI SUBCUTANEOUS SOLUTION AUTO-INJECTOR .....
RYDAPT.....	12, 13	168, 169
RYKINDO .....	160	SIMPONI SUBCUTANEOUS SOLUTION PREFILLED SYRINGE .....
RYLAZE.....	161	168, 169
<b>S</b>		sirolimus oral solution 1 mg/ml .....
sapropterin dihydrochloride oral packet .	162	237
		sirolimus oral tablet 0.5 mg, 1 mg, 2 mg
		237
		SIRTURO .....
		170
		SKYTROFA .....
		77
		sodium oxybate .....
		171
		sodium phenylbutyrate oral powder 3 gm/tsp .....
		172
		sodium phenylbutyrate oral tablet.....
		172
		SOFOSBUVIR-VELPATASVIR.....
		173
		SOLTAMOX .....
		12, 13
		SOMAVERT.....
		174
		sorafenib tosylate .....
		12, 13
		SOTYKTU .....
		175
		STARJEMZA INTRAVENOUS ....
		202, 203
		STARJEMZA SUBCUTANEOUS SOLUTION.....
		202, 203

STARJEMZA SUBCUTANEOUS  
 SOLUTION PREFILLED SYRINGE 45  
 MG/0.5ML, 90 MG/ML ..... 202, 203  
 STELARA INTRAVENOUS ..... 202, 203  
 STELARA SUBCUTANEOUS  
 SOLUTION 45 MG/0.5ML ..... 202, 203  
 STELARA SUBCUTANEOUS  
 SOLUTION PREFILLED SYRINGE 45  
 MG/0.5ML, 90 MG/ML ..... 202, 203  
 STEQEYMA INTRAVENOUS ..... 202, 203  
 STEQEYMA SUBCUTANEOUS  
 SOLUTION..... 202, 203  
 STEQEYMA SUBCUTANEOUS  
 SOLUTION PREFILLED SYRINGE 45  
 MG/0.5ML, 90 MG/ML ..... 202, 203  
 STIVARGA ..... 12, 13  
 SUCRAID ..... 176  
 sunitinib malate..... 12, 13  
 SYMDEKO..... 177  
 SYNAREL ..... 178  
**T**  
 TABLOID ..... 12, 13  
 TABRECTA ..... 12, 13  
 tacrolimus intravenous solution 5 mg/ml 237  
 tacrolimus oral capsule 0.5 mg, 1 mg, 5 mg  
 ..... 237  
 tadalafil (pah)..... 179  
 tadalafil oral tablet 5 mg ..... 180  
 TADLIQ..... 179  
 TAFINLAR..... 12, 13  
 TAGRISSO ..... 12, 13  
 TALTZ SUBCUTANEOUS SOLUTION  
 AUTO-INJECTOR ..... 181, 182  
 TALTZ SUBCUTANEOUS SOLUTION  
 PREFILLED SYRINGE 20 MG/0.25ML,  
 40 MG/0.5ML, 80 MG/ML ..... 181, 182  
 TALZENNA ..... 12, 13  
 TARPEYO ..... 183  
 TASCENSO ODT..... 114, 115  
 tasimelteon ..... 184  
 TAVNEOS..... 185, 186  
 TAZVERIK..... 12, 13  
 temazepam ..... 85  
 TENIVAC INTRAMUSCULAR  
 INJECTABLE 5-2 LFU (INJECTION)  
 ..... 237

TENIVAC INTRAMUSCULAR  
 SUSPENSION 5-2 LF/0.5ML ..... 237  
 TEPEZZA ..... 187  
 TEPMETKO ..... 12, 13  
 teriflunomide..... 114, 115  
 teriparatide subcutaneous solution pen-  
 injector 560 mcg/2.24ml ..... 188  
 testosterone transdermal gel 1.62 %, 12.5  
 mg/act (1%), 20.25 mg/1.25gm (1.62%),  
 20.25 mg/act (1.62%), 25 mg/2.5gm  
 (1%), 40.5 mg/2.5gm (1.62%), 50  
 mg/5gm (1%) ..... 194  
 testosterone transdermal solution..... 194  
 tetrabenazine ..... 210  
 THALOMID ORAL CAPSULE 100 MG,  
 50 MG ..... 12, 13  
 TIBSOVO ..... 12, 13  
 tigecycline..... 190  
 tiopronin oral..... 189  
 TOBI PODHALER..... 191  
 tobramycin inhalation nebulization solution  
 300 mg/4ml, 300 mg/5ml..... 237  
 tolvaptan..... 192  
 tolvaptan (hyponatremia)..... 192  
 topiramate oral solution ..... 60  
 toremifene citrate ..... 12, 13  
 TRELSTAR MIXJECT ..... 75  
 TREMFYA ONE-PRESS  
 SUBCUTANEOUS SOLUTION PEN-  
 INJECTOR..... 196  
 TREMFYA PEN SUBCUTANEOUS  
 SOLUTION AUTO-INJECTOR 100  
 MG/ML, 200 MG/2ML ..... 196  
 TREMFYA SUBCUTANEOUS  
 SOLUTION PREFILLED SYRINGE 100  
 MG/ML, 200 MG/2ML ..... 196  
 TREMFYA-CD/UC INDUCTION..... 196  
 tretinoin oral..... 12, 13  
 trientine hcl oral capsule 250 mg..... 197  
 TRIKAFTA..... 198  
 TRULICITY SUBCUTANEOUS  
 SOLUTION AUTO-INJECTOR ..... 74  
 TRUQAP ORAL TABLET 200 MG.. 12, 13  
 TRUQAP ORAL TABLET THERAPY  
 PACK..... 12, 13  
 TUKYSA ..... 12, 13

TURALIO ORAL CAPSULE 125 MG... 12, 13  
 TYMLOS ..... 199  
 TYVASO DPI MAINTENANCE KIT  
 INHALATION POWDER 112 X 32MCG & 112 X64MCG, 112 X 48MCG & 112 X64MCG, 16 MCG, 32 MCG, 48 MCG, 64 MCG, 80 MCG ..... 200  
 TYVASO DPI TITRATION KIT  
 INHALATION POWDER 16 & 32 & 48 MCG ..... 200  
**U**  
 UBRELVY ..... 32, 33  
 UPTRAVI ORAL ..... 201  
 UPTRAVI TITRATION..... 201  
 ustekinumab subcutaneous solution 202, 203  
 ustekinumab subcutaneous solution prefilled syringe 45 mg/0.5ml, 90 mg/ml.. 202, 203  
 ustekinumab-aauz subcutaneous solution prefilled syringe 45 mg/0.5ml, 90 mg/ml ..... 202, 203  
 ustekinumab-aekn subcutaneous solution prefilled syringe 45 mg/0.5ml, 90 mg/ml ..... 202, 203  
 UZEDY SUBCUTANEOUS SUSPENSION  
 PREFILLED SYRINGE 100 MG/0.28ML, 125 MG/0.35ML, 150 MG/0.42ML, 200 MG/0.56ML, 250 MG/0.7ML, 50 MG/0.14ML, 75 MG/0.21ML ..... 204  
**V**  
 VALCHLOR..... 205  
 VALTOCO 10 MG DOSE..... 118  
 VALTOCO 15 MG DOSE NASAL LIQUID THERAPY PACK 2 X 7.5 MG/0.1ML ..... 118  
 VALTOCO 20 MG DOSE NASAL LIQUID THERAPY PACK 2 X 10 MG/0.1ML ..... 118  
 VALTOCO 5 MG DOSE..... 118  
 VANFLYTA ..... 12, 13  
 VEMLIDY ..... 206  
 VENCLEXTA..... 12, 13  
 VENCLEXTA STARTING PACK ... 12, 13  
 VEOZAH ..... 207  
 VERZENIO..... 12, 13

vigabatrin ..... 208  
 VIGAFYDE ..... 208  
 VIJOICE ..... 209  
 VITRAKVI ..... 12, 13  
 VIZIMPRO ..... 12, 13  
 VONJO ..... 12, 13  
 VORANIGO ..... 12, 13  
 voriconazole intravenous ..... 211  
 VOSEVI..... 212  
 VOWST ..... 213  
 VRAYLAR ORAL CAPSULE..... 128, 129  
**W**  
 WEGOVY ORAL ..... 216, 217  
 WEGOVY SUBCUTANEOUS SOLUTION AUTO-INJECTOR 0.25 MG/0.5ML, 0.5 MG/0.5ML, 1 MG/0.5ML, 1.7 MG/0.75ML, 2.4 MG/0.75ML ... 214, 215  
 WELIREG..... 12, 13  
 WINREVAIR..... 219  
**X**  
 XALKORI..... 12, 13  
 XATMEP ..... 109  
 XDEMVY ..... 220  
 XELJANZ ORAL SOLUTION ..... 221  
 XELJANZ ORAL TABLET..... 221  
 XELJANZ XR ..... 221  
 XERMELO ..... 222  
 XIFAXAN..... 223  
 XOLAIR ..... 224, 225  
 XOLREMDI ..... 226  
 XOSPATA ..... 12, 13  
 XPOVIO (100 MG ONCE WEEKLY) ORAL TABLET THERAPY PACK 50 MG ..... 12, 13  
 XPOVIO (40 MG ONCE WEEKLY) ORAL TABLET THERAPY PACK 10 MG, 40 MG ..... 12, 13  
 XPOVIO (40 MG TWICE WEEKLY) ORAL TABLET THERAPY PACK 40 MG ..... 12, 13  
 XPOVIO (60 MG ONCE WEEKLY) ORAL TABLET THERAPY PACK 60 MG... 12, 13  
 XPOVIO (60 MG TWICE WEEKLY)12, 13

XPOVIO (80 MG ONCE WEEKLY) ORAL  
 TABLET THERAPY PACK 40 MG, 80  
 MG ..... 12, 13  
 XPOVIO (80 MG TWICE WEEKLY) 12, 13  
 XTANDI ..... 12, 13  
 XYWAV ..... 227  
**Y**  
 YESINTEK INTRAVENOUS ..... 202, 203  
 YESINTEK SUBCUTANEOUS  
 SOLUTION..... 202, 203  
 YESINTEK SUBCUTANEOUS  
 SOLUTION PREFILLED SYRINGE 45  
 MG/0.5ML, 90 MG/ML ..... 202, 203  
 YONSA..... 12, 13  
 YORVIPATH ..... 228  
 YUTREPIA..... 229  
**Z**  
 zaleplon oral capsule 10 mg, 5 mg..... 85  
 ZARXIO ..... 218  
 ZAVZPRET ..... 32, 33

ZEJULA ORAL TABLET..... 12, 13  
 ZELBORAF ..... 12, 13  
 ZEMAIRA ..... 8  
 ZEPBOUND SUBCUTANEOUS  
 SOLUTION AUTO-INJECTOR 230, 231  
 ZEPOSIA ..... 232  
 ZEPOSIA 7-DAY STARTER PACK..... 232  
 ZEPOSIA STARTER KIT ORAL  
 CAPSULE THERAPY PACK 0.23MG  
 &0.46MG 0.92MG(21)..... 232  
 ZILBRYSQ ..... 233  
 ZOLINZA ..... 12, 13  
 zolpidem tartrate er ..... 85  
 zolpidem tartrate oral tablet 10 mg ..... 85  
 ZTALMY ..... 234  
 ZTLIDO ..... 195  
 ZURZUVAE ..... 235  
 ZYDELIG ..... 12, 13  
 ZYKADIA ORAL TABLET ..... 12, 13